

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁵ : C12N 15/53, 15/82, 5/10 A01H 5/00, 5/10, C12N 9/02</p>	<p>A1</p>	<p>(11) International Publication Number: WO 90/02804 (43) International Publication Date: 22 March 1990 (22.03.90)</p>
<p>(21) International Application Number: PCT/EP89/01025 (22) International Filing Date: 31 August 1989 (31.08.89) (30) Priority data: 88402222.9 2 September 1988 (02.09.88) EP (34) Countries for which the regional or international application was filed: AT et al. (71) Applicant (for all designated States except US): PLANT GENETIC SYSTEMS, N.V. [BE/BE]; Kolonel Bourg- straat 106, B-1040 Brussels (BE). (72) Inventors; and (75) Inventors/Applicants (for US only) : BOWLER, Chris [GB/ BE]; Parklaan 13, B-9000 Gent (BE). INZE, Dirk [BE/ BE]; Driesstraat 18, B-9390 Moorsel-Aalst (BE). VAN CAMP, Wim [BE/BE]; Tulpenlaan 21, B-3540 Zolder (BE). ALLIOTTE, Thierry [FR/BE]; Kortrijksesteenweg 989, B-9000 Gent (BE). VAN MONTAGU, Marc [BE/ BE]; De Stassartstraat 120, B-1050 Brussels (BE). BOT- TERMAN, Johan [BE/BE]; Het Wijngaardeke 5,</p>		<p>B-9721 Zevergem-De Pinte (BE). (74) Agent: GUTMANN, Ernest; SC Ernest Gutmann-Yves Plasseraud, 67, boulevard Haussmann, F-75008 Paris (FR). (81) Designated States: AU, JP, US. Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>
<p>(54) Title: STRESS-TOLERANT PLANTS</p> <p>(57) Abstract</p> <p>A plant, the nuclear genome of which is transformed with a recombinant DNA sequence encoding a superoxide dismutase which renders the plant stress-resistant. The recombinant DNA sequence also optionally encodes a targeting peptide, fused to the superoxide dismutase, so that the superoxide dismutase is expressed in the cytoplasm of the plant's cells and is subsequently targeted to mitochondria or chloroplasts of the plant's cells or is secreted, via the endoplasmic reticulum, from the plant's cells.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauntania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Fasso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LJ	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

STRESS-TOLERANT PLANTS

5 This invention relates to DNA molecules and genes coding for metallo-superoxide dismutase enzymes (the "SODs"), particularly plant SODs, particularly a plant manganese superoxide dismutase (the "MnSOD") or a plant iron superoxide dismutase (the "FeSOD"). Production, particularly overproduction, of one or more SOD enzymes can be used to confer on a plant resistance or tolerance
10 to toxic, highly reactive, oxygen species, particularly superoxide anions, produced in the plant's cells under many naturally occurring stress conditions.

This invention also relates to a recombinant gene (the "recombinant SOD gene") which is preferably a chimaeric
15 gene and which contains the following operably linked DNA fragments in the same transcriptional unit: 1) a DNA sequence encoding an SOD (the "SOD gene"), preferably a DNA sequence encoding MnSOD or FeSOD (the "MnSOD gene" or "FeSOD gene", respectively); 2) a promoter suitable for
20 controlling transcription of the SOD gene in a plant cell; and 3) suitable transcription termination and polyadenylation signals for expressing the SOD gene in a plant cell. The recombinant SOD gene optionally contains an additional DNA fragment encoding a targeting peptide
25 (the "targeting sequence") immediately upstream of, and in the same reading frame as, the SOD gene, whereby a plant cell, transformed with the recombinant SOD gene, produces or overproduces a precursor of the SOD having an N-terminal peptide characteristic for: 1) mitochondrial or
30 chloroplast targeting of the SOD within the plant cell; or 2) translocation of the SOD to the lumen of the endoplasmatic reticulum ("ER") of the plant cell for eventual secretion of the SOD out of the plant cell.

This invention further relates to the use of an SOD gene, particularly a recombinant SOD gene, in the production of a transgenic plant having an increased resistance or tolerance to stress conditions which produce highly reactive oxygen species, particularly superoxide anions, in one or more compartments of the plant's cells. This invention relates particularly to the use of the SOD gene for the protection of the plant against naturally occurring stress conditions which are not normally within the control of a farmer (e.g., conditions of soil composition, climate, etc). As a result, the invention provides a means for growing crops in geographical areas in which they could not heretofore be grown with reasonable yields due to such naturally occurring stress conditions.

This invention still further relates to: a cell of a plant (the "transgenic plant cell"), the genome, particularly the nuclear genome, of which is transformed with the recombinant SOD gene; a culture of such cells; a plant (the "transgenic plant") which is regenerated from the transgenic plant cell or is produced from a so-regenerated plant and the genome of which contains the recombinant SOD gene; and the reproductive materials (e.g., seeds) of the transgenic plant. The transgenic plant is resistant or tolerant to stress conditions, particularly naturally occurring stress conditions, which produce highly reactive oxygen species in one or more compartments of the plant's cells, thereby increasing the potential yield and/or quality of crops produced by the plant.

BACKGROUND OF THE INVENTION

Plants have to be able to cope with a large number of naturally occurring physicochemical stress situations such


as drought, waterlogging, high salt concentrations, high or low temperatures, metal excess and metal starvation, as well as biological stress situations such as various pathogens. These stress situations can interfere with normal plant growth and development and consequently, in the case of crop plants, can lower food quality and yield.

It is already known that the production of certain proteins is induced in plants by different stress conditions (Sachs and Ho, 1986). The characteristics of the induced proteins depend upon the actual toxicity effects produced by the stress conditions in the plants. Recently, toxicity due to highly reactive oxygen species has been recognized as an important component of the deleterious effects of a number of stress conditions on plants.

Highly reactive oxygen species are generated in plant cells by the action of various agents such as herbicides (Harbour and Bolton, 1975; Orr and Hogan, 1983), air pollutants (Grimes et al., 1983; Tanaka et al., 1982) such as ozone (MacKay et al., 1987), redox active compounds (Hassan and Fridovich, 1979), heat shock (Lee et al., 1983) and chilling (Clare et al., 1984), and they can cause biologically significant cellular injury. Indeed, superoxide species (O_2^-), hydrogen peroxide (H_2O_2) and the hydroxyl radical (OH^-) can initiate peroxidation of membrane lipids (Mead, 1976), mark proteins for proteolysis (Fucci et al., 1983), cause DNA damage (Brown and Fridovich, 1981; Imlay and Linn, 1988), inhibit photosynthesis (Robinson et al., 1980; Kaiser, 1979) and destroy chlorophyll (Harbour and Bolton, 1978).

Oxygen radicals are also produced under normal conditions in chloroplasts under illumination (Asada et al., 1974), metabolically as products of enzymes (Fridovich, 1978) and in beta-oxidation of fatty acids

(Beavers, 1979). All organisms have developed ways to destroy these toxic, highly reactive oxygen species produced under normal conditions. Generally, the superoxide anion is converted to hydrogen peroxide by the action of an SOD in the following reaction:



Catalase then catalyzes the decomposition of 2 moles of H_2O_2 into 2 moles of H_2O and 1 mole of O_2 and thus protects the cells against the noxious H_2O_2 . In plant cells, peroxidases also provide an important alternative pathway to eliminate H_2O_2 . SOD and catalase are ubiquitous in aerobic prokaryotic and eukaryotic cells.

SODs are a group of metalloproteins which have been classified according to their metal cofactor (Bannister et al., 1987). Iron enzymes (FeSODs) are present in some prokaryotes and occasionally in plants. Manganese enzymes (MnSODs) are widely distributed among prokaryotic and eukaryotic organisms, and in eukaryotes, they are most often found in the mitochondrial matrix. Copper-zinc enzymes (Cu/ZnSODs) are found almost exclusively in eukaryotes where they are often present in several isoforms. SODs are produced by plant cells as part of their natural defense mechanisms against the toxic effects of highly reactive oxygen species.

The isolation of several cDNA's and genes encoding SODs from various species has been reported (Parker and Blake; 1988; Ho and Crapo, 1988; Birmingham-McDonogh et al., 1988; Marres et al., 1985; Seto et al., 1987; Carliz et al., 1988). In plants, little is known at the molecular level about SODs. A cDNA clone encoding a cytosolic Cu/ZnSOD has been isolated from maize (Cannon et al., 1987), and a cDNA clone encoding MnSOD from *N.*

plumbaginifolia has been reported (Bauw et al., 1987). However, no sequence data have been published for plant MnSOD or FeSOD genes.

It is known that increased oxygen radicals, produced in plants under oxidative stress conditions, influence the levels of oxygen radical-detoxifying enzymes such as SODs. Resistance against oxygen radical-producing herbicides, such as paraquat (i.e., methyl viologen), has been shown to be correlated with increased levels of enzymes involved in superoxide detoxification in Conyza spec. (Shaaltiel and Gressel, 1987). Similar results have been obtained in Chlorella sorokiniana (Rabinowitch and Fridovich, 1985). Increased Cu/ZnSOD has also been observed in paraquat resistant calluses of Nicotiana tabacum (Furasawa et al., 1984). Paraquat also induces 40% more Cu/ZnSOD in maize whereas only a negligible induction of MnSOD is observed (Matters and Scandalios, 1986). Australian patent AU-A-27461/84 discloses that: paraquat can be used as an efficient weed-killer in postemergence stages, provided that crop plants are made tolerant to its toxic effect; a cDNA clone for human Cu/ZnSOD can be used to identify and isolate a plant DNA segment which carries a plant gene encoding Cu/ZnSOD; and the plant gene encoding Cu/ZnSOD can be inserted, using known expression vectors, into plant cells in order to make paraquat-resistant plants.

The deleterious effects of some air pollutants also seem to be mediated through oxygen radicals. Young poplar leaves, which contain 5 times more SOD than old leaves, are more resistant to SO₂ toxicity (Tanaka and Sugahara, 1980). Addition of N-(2-(2-oxo-1-imidazolidenyl)ethyl)-N'-phenylurea ("EDU"), an antiozonant, reduces injury to a plant from ozone. This correlates with an increase in SOD activity in Phaseolus vulgaris (Lee and Bennett, 1982). DDR patent publication 225716 describes the detection of

pollution resistant plants by measuring their Cu/ZnSOD content as an indicator of SO₂ tolerance.

Green plants are subject to more or less severe damage by combinations of high light intensities with either high or low temperatures. The elevated production of oxygen radicals under these conditions has been implicated as a cause of so-called photooxidative or photodynamic damage. SOD levels in plants are inversely correlated with the extent of the incurred photooxidative damage. This has been reported in ripening tomatoes in which susceptibility to sunscald (a special kind of photodynamic damage) was directly related to SOD activity both under natural and experimental conditions (Rabinowith et al., 1982).

Increased resistance to chilling injury in Chlorella ellipsoidea and to photooxidative death in Plectonema boryanum has also been correlated with SOD activity (Steinitz et al., 1979; Clare et al., 1984).

SOD activity has also been connected to tolerance to hyperoxic or anoxic conditions. The presence of the enzyme in rhizomes of Iris pseudoacorus under anaerobic conditions was seen to be important for its ability to recover from anoxic stress, and the efficacy of recovery was correlated with the SOD level (Monk et al., 1987).

An important naturally occurring stress condition for most plants is the presence of various pathogens which induce so-called "pathogenesis related" ("PR") proteins (Sachs and Ho, 1986; Collinge and Slusarenko, 1987). Major changes in SOD activity apparently take place when plants are attacked by pathogens. Increases in enzymatic activity have been found in susceptible plants while decreases were noticed in resistant ones. Examples are pea roots infected by the cyst nematode Heterodera goettingiana (Arrigoni et al., 1981) and tomato plants infested with Meloidogyne incognita (Zacheo et al., 1982). Plant parasitic fungi of

the genera Alternaria and Cercospora are known to produce toxins (altertoxins and cercosporins, respectively) which generate singlet oxygen and superoxide (Daub, 1982, 1987; Daub and Briggs, 1983; Daub and Hangarter, 1983; Hartman et al., 1989). Tobacco plants regenerated from calli selected for high SOD activity were shown to be resistant to cercosporin (Furusawa and Mizuguchi, 1983).

It is also believed possible that SOD is active in the defense of plants against damage from ionizing radiation. At least in the fruit fly Drosophila, a natural genetic polymorphism has been found in regard to SOD. Flies with the greatest resistance against ionizing radiation carry an allele which codes for an SOD enzyme with a higher specific activity (Peng et al., 1986).

Reduced oxygen species also appear to play a key role in the aging process of plants (Munkres et al., 1984).

SUMMARY OF THE INVENTION

In accordance with this invention, a plant MnSOD gene and FeSOD gene and the MnSOD and FeSOD, which can be produced by expression of these genes in procaryotic or eucaryotic cells, particularly plant cells, are provided.

Also in accordance with this invention, a recombinant SOD gene, preferably a chimaeric recombinant SOD gene, is provided for stably transforming a plant cell genome, containing the following operably linked DNA fragments: 1) a SOD gene; 2) a promotor capable of directing expression of the SOD gene in a plant cell; and 3) suitable 3' transcription regulation signals. Preferably, the recombinant SOD gene also contains a targeting sequence which is located between the promoter and the SOD gene, which is in the same reading frame as, and fused to, the SOD gene, and which codes for a mitochondrial or chloroplastic targeting peptide of the plant cell or for a

targeting peptide for translocation to the lumen of the ER of the plant cell, so that a fusion protein containing the targeting peptide and the SOD can be synthesized in the cytoplasm of the plant cell and so that the fusion protein will be processed by the plant cell whereby the SOD is translocated into mitochondria or chloroplasts of the plant cell or into the lumen of the ER for secretion out of the plant cell.

Further in accordance with this invention are provided transgenic plant cells, cultures of the plant cells, and transgenic stress-resistant plants regenerated from the cells, which have stably incorporated into their genome, preferably their nuclear genome, one or more of the recombinant SOD genes, so that at least one SOD is secreted from the cells or is expressed or overexpressed in the cytosol, the mitochondria and/or the chloroplasts of the cells.

DESCRIPTION OF THE INVENTION

The plant MnSOD gene of the present invention can be identified, isolated and characterized using methods well known to those skilled in the art. The aminoterminal sequence of proteins electroblotted on support material after two dimensional gel electrophoresis can be determined in a straightforward manner by gas-phase sequencing of the immobilized proteins (Bauw et al., 1987). In this way the amino-terminal aminoacid sequence of the MnSOD of Nicotiana plumbaginifolia was determined. The construction of an oligonucleotide probe specific for this aminoacid sequence then allows the isolation of the cDNA encoding the MnSOD from a cDNA library of Nicotiana plumbaginifolia. The cDNA can subsequently be manipulated and sequenced using standard methods (Maniatis et al., 1982; Maxam and Gilbert, 1980). The complete aminoacid

sequence of the plant MnSOD can then be deduced from the cDNA sequence. It goes without saying that, not only the specific cDNA sequences of Figure 1, 2 and 3, but also all DNA sequences coding for proteins with the deduced amino acid sequences of Figure 1, 2, and 3 and their equivalents with MnSOD activity, fall within the scope of this invention.

The isolated cDNA of the MnSOD gene contains not only the complete reading frame of 204 aminoacids comprising the mature active MnSOD but also a transit peptide-encoding sequence of 24 aminoacids which targets the enzyme into the mitochondria as deduced from the general properties of mitochondrial transit peptides (Schatz, 1987). The complete nucleotide and amino acid sequence of the cDNA coding for the mitochondrial transit peptide and MnSOD is shown in Fig. 1.

The available data on plant FeSOD genes suggest that they are fairly different from procaryotic FeSOD genes. However, plant FeSOD genes can be identified by complementation of a SOD deficient E. coli strain. In this procedure comprised of individual steps well known to those skilled in the art, a cDNA library of Nicotiana plumbaginifolia can be cloned in an E. coli expression vector, such as pUC18 (Yannisch-Perron et al., 1985), and the resulting plasmids can be used to transform a SOD deficient E. coli strain such as that described by Carlroz and Touati (1987) by electroporation in order to obtain high transformation efficiency. Colonies that are able to grow aerobically on minimal medium and that are synthesizing SOD can be identified by staining for SOD activity of total cellular protein separated by polyacrylamide gel electrophoresis. The cDNA inserts in the expression vector can then be further characterized by restriction analysis and/or hybridization studies. Inserts

10

that are thought to be coding for a FeSOD (i.e., that do not hybridize with known plant Cu/ZnSODs and MnSOD and that direct the production of an active SOD that is resistant to KCN and sensitive to H_2O_2) are selected for further characterization by means of DNA sequencing (Sanger et al., 1977). The aminoacid sequence of the FeSOD can then be inferred from the obtained nucleotide sequence. Using this procedure a cDNA coding for a FeSOD from Nicotiana plumbaginifolia, characterized by the sequence in Fig. 5, could be obtained.

Although the plant SOD genes, particularly the plant MnSOD and FeSOD genes, are the preferred SOD genes in the recombinant SOD genes of this invention, other genes coding for enzymes with SOD activity, such as procaryotic SOD genes and cDNAs derived from eucaryotic SOD genes, also can be used. In this regard, the selection of the SOD gene is not believed to be critical, and a particular plant cell can be transformed with a recombinant SOD gene containing either: a foreign SOD gene which will produce its encoded SOD in the cell; or an endogenous SOD gene which will provide overproduction of its encoded SOD in the cell. For example, suitable foreign SOD genes can encode: the procaryotic and eucaryotic Cu/ZnSODs listed by Getzoff et al. (1989); the procaryotic and eucaryotic MnSODs listed in Bowler et al. (1989a); and the procaryotic FeSODs described by Carlizoz et al., 1988) and Parker and Blake (1988).

For constructing a recombinant SOD gene which can be expressed in a transformed plant cell, suitable promoters are known which can be provided upstream (i.e., 5') of a SOD gene, and the selection of a promoter is not believed to be critical. In this regard, a particular plant cell can be transformed with a recombinant SOD gene containing either a foreign or an endogenous promoter suitable for

directing expression or overexpression of the SOD gene. Suitable foreign constitutive promoters include: the promoter of the cauliflower mosaic virus ("CaMV") isolates CM1841 (Gardner et al., 1981) and CabbB-S (Franck et al., 1980) [the "35S promoter"] which directs constitutive expression of heterologous genes (Odell et al., 1983); a related promoter (the "35S3 promoter") which can be isolated from the CaMV isolate CabbB-JI (Hull and Howell, 1978) and which differs from the 35S promoter in sequence and in its greater activity in transgenic plants (Harpster et al., 1988); and the TR1' and the TR2' promoters which drive the expression of the 1' and 2' genes, respectively, of the T-DNA (Velten et al., 1984) and are wound-induced promoters. Suitable organ-specific, tissue-specific and/or inducible foreign promoters are also known (Kuhlemeier et al., 1987 and the references cited therein) such as the promoter of the 1A small subunit gene of 1,5 ribulose biphosphate carboxylase (Rubisco) of Arabidopsis thaliana (the "ssu promoter") which is a light-inducible promoter (European patent 242,246) active only in photosynthetic tissue. Other organ-specific, tissue-specific and inducible promoters can be isolated from cell- or tissue- or organ-specific genes and from genes specific for particular developmental stages (Goldberg, 1988) by the screening of plant genomic libraries with specific cDNAs, using techniques as disclosed, for example, in European patent applications 89402224.3 and 89401194.9.

For constructing a recombinant SOD gene which can be expressed in a transformed plant cell, suitable transcription termination and polyadenylation signals are known which can be provided downstream (i.e., 3') of an SOD gene, and the selection of such 3' transcription regulation signals is not critical. In this regard, a particular plant cell can be transformed with a

12

recombinant SOD gene containing either foreign or endogeneous transcription termination and polyadenylation signals suitable for obtaining expression or overexpression of the SOD gene. For example, the foreign 3' untranslated ends of genes, such as gene 7 (Velten and Schell, 1985), the octopine synthase gene (Gielen et al., 1983) and the nopaline synthase gene of the T-DNA region of Agrobacterium tumefaciens Ti-plasmid (European patent application 89402224.3), can be used.

By "recombinant" with regard to the recombinant SOD gene of this invention is meant that its operably linked SOD gene, promoter and 3' transcription regulation signals, together with any targeting sequence, can be introduced in a plant genome by artificial means (e.g., by Agrobacterium-mediated gene transfer) and are then (in the plant genome) not in their natural genomic environment (i.e., are not surrounded in the plant genome by their naturally surrounding DNA sequences).

By "chimaeric" with regard to the chimaeric recombinant SOD gene of this invention is meant that its SOD gene: 1) is not naturally found under the control of its promoter and/or 2) is not naturally found fused to, and in the same reading frame as, its targeting sequence. Examples of chimaeric recombinant SOD genes of this invention comprise: an SOD gene of bacterial origin under the control of a promoter of plant origin; and a SOD gene of plant origin under the control of a promoter of viral origin and fused to a signal sequence encoding a transit peptide of plant origin.

For constructing a recombinant SOD gene which can be expressed in a transformed plant cell, preferably in its cytoplasm, by the production or overproduction of an SOD, followed by translocation to the cell's mitochondria, chloroplasts and/or lumen of the cell's endoplasmatic

reticulum ("ER") for eventual secretion from the cell, suitable targeting sequences encoding targeting peptides are known. Selection of such targeting sequences is not believed to be critical, and a particular plant cell can be transformed with a recombinant SOD gene containing either a foreign or an endogeneous targeting sequence which will provide translocation or secretion of the expression product of the SOD gene.

By "targeting peptide" is meant a polypeptide fragment which is normally associated in a eucaryotic cell with a chloroplast or mitochondrial protein or subunit of the protein or with a protein translocated to the ER and which is produced in a cell as part of a precursor protein encoded by the nuclear DNA of the cell. The targeting peptide is responsible for the translocation process of the nuclear-encoded chloroplast or mitochondrial protein or subunit into the chloroplast or the mitochondria or the lumen of the ER. During the translocation process, the targeting peptide is separated or proteolytically removed from the protein or subunit. A targeting sequence can be provided in the recombinant SOD gene of this invention for providing a targeting peptide to translocate an expressed SOD within a transformed plant cell as generally described in European patent applications 85402596.2 and 88402222.9. A suitable targeting peptide for transport into chloroplasts is the transit peptide of the small subunit of the enzyme RUBP carboxylase (European patent application 85402596.2), but other chloroplast transit peptides, such as those listed by Watson (1984), can also be used. A suitable mitochondrial targeting peptide is the transit peptide naturally associated with a plant MnSOD as shown in Figure 1, but other mitochondrial transit peptides, such as those described by Schatz (1987) and listed by Watson (1984), can be used. Suitable targeting

peptides that can translocate an SOD to the lumen of the ER of a plant cell are, for instance, the signal peptides described by Von Heijne (1988) and listed by Watson (1984).

5 In accordance with this invention, any SOD gene, preferably a plant SOD gene, particularly a plant MnSOD or FeSOD gene, can be used to produce a transformed plant with increased tolerance or resistance to the toxic effects of highly reactive oxygen species, particularly
10 superoxide anions, produced in one or more compartments of its cells as a result of certain stress conditions, oxidation during or after harvest, or senescence. Preferably, the resulting SOD is produced in, or transported to, mitochondria, chloroplasts, cytosols or
15 other specific sites in or out the plant cells, appropriate to combatting the effects of the toxic highly reactive oxygen species. In this regard, the targeting sequence encoding a targeting peptide in the recombinant SOD gene permits the SOD expression product to be targeted
20 to one or more cell compartments in or out of the plant cells where stress-produced, highly reactive, oxygen species pose a particular problem as discussed below.

The recombinant SOD gene of this invention, which is preferably chimaeric, can be used to produce transgenic
25 plant cells and transgenic plants in which the SOD gene is expressed or preferably overexpressed. This confers on these cells and plants an increased resistance to the toxicity of highly reactive oxygen species, particularly superoxide radicals. Since the formation of highly
30 reactive oxygen species is likely to occur at different sites within the plant cells, depending upon the nature of the stress to which the cells are subjected, it is important to be able to direct expression of the SOD gene into one or more compartments of the cells containing such
35

5 sites. Directing SOD gene overexpression into chloroplasts can be important for protection against stress induced by conditions such as high light intensities in combination with high or low temperatures (Bowles, 1984) or the
10 presence of paraquat (Shaaltiel and Gressel, 1987), which generate highly reactive oxygen species in these organelles. The targeting of SOD overexpression into mitochondria aims at protecting plants against deleterious effects of highly reactive oxygen species generated in
15 these organelles, for example by plant pathogens which can cause increased respirational activity in mitochondria. SOD overexpression in the cytosol can protect plants against the effects of highly reactive oxygen species generated in this cell compartment, thereby leading, for example, to increased storage life of the plants and to their fruit having increased tolerance to bruising. SOD
20 secretion from plant cells can provide increased resistance to reactive oxygen species outside the plant cells, such as are caused by infections of pathogens responsible for the production of highly reactive oxygen species (e.g., by the fungus Cercospora) and by air pollutants (e.g., ozone and SO₂). It is also believed that SOD-overexpressing plants may have a prolonged lifespan due to their higher level of protection against
25 toxic oxygen species.

A transgenic plant of this invention can be produced by the introduction of the recombinant SOD gene into a cell of the plant, followed by regeneration of the plant, using known techniques. A disarmed Agrobacterium tumefaciens
30 Ti-plasmid can be used for transforming the plant cell with the recombinant SOD gene using procedures described, for example, in European patent publications 116718 and 270822, PCT publication WO 84/02913 and European patent application 87400544.0. Preferably, the
35 Ti-plasmid contains the recombinant SOD gene between its

16

border sequences or at least located to the left of the right border sequence of the T-DNA of the Ti-plasmid. Such a Ti-plasmid can be used for the transformation of all plants susceptible to Agrobacterium infection, such as Nicotiana plumbaginifolia, Arabidopsis thaliana Nicotiana tabacum, Solanum tuberosum, Lycopersicum esculentum, Medicago sativa and Beta vulgaris. Other techniques can be used to transform these and other plants, such as: direct gene transfer (as described, for example, in European patent publication 223247), pollen mediated transformation (as described, for example, in European patent publication 270356, PCT publication WO 85/01856 and European patent publication 275069), in vitro protoplast transformation (as described, for example, in US patent 4684611), plant RNA virus-mediated transformation (as described, for example, in European patent publication 67553 and US patent 4407956) and liposome-mediated transformation (as described, for example, in US patent 4536475).

The resulting transgenic plant can be used in a conventional plant breeding scheme to produce more transgenic plants with the same characteristics or to introduce the recombinant SOD gene in other varieties of the same or related plant species. Reproductive materials (e.g., seeds), which are obtained from the transgenic plants, contain the recombinant SOD gene as a stable genomic insert.

The production of transgenic plants, having two or more recombinant SOD genes which are stably integrated in their genomes, particularly their nuclear genomes, and which target the expression or overexpression of one or more SODs into several different compartments of the plants' cells at once (e.g., with targeting sequences encoding different targeting peptides), can provide additional benefits in stress tolerance or resistance.

Such a stress-resistant plant can be obtained in several ways, such as by: several independent transformations of the same plant with different recombinant SOD genes; a single transformation with a vector containing two or more recombinant SOD genes in tandem; or crossing two plants, each of which has already been transformed with different recombinant SOD genes. In order to be able to adequately select plants with two recombinant SOD genes, it is advisable that each recombinant SOD gene contain, within the same genetic locus, a different selectable marker gene. Selectable marker genes that can be used for this purpose are, for instance, the neo gene coding for neomycin phosphotransferase (Reiss et al., 1984) and the bar gene coding for phosphinotricin acetyltransferase (as described in European patent publication 242246).

The transgenic plant cells, cell cultures and plants of this invention can also be used to produce an overexpressed SOD, especially a plant SOD, particularly a plant MnSOD or FeSOD. In a conventional manner, the so-produced SOD can be recovered from the plant cells and used as an anti-oxidative food additive, an anti-inflammatory agent in mammals, or a therapeutic agent in mammals for certain pathological conditions that generate superoxide radicals or for prevention of ischemic injuries (See Bannister et al., 1987).

An SOD of this invention can also be produced, preferably as a secreted protein, by linking an SOD gene to suitable expression signals such as appropriate promoter sequences, secretion targeting sequences, ribosome binding sites, start and stop codons and transcription regulation sequences for prokaryotic and/or eukaryotic cells. The so-linked SOD gene can then be introduced into a host, such as a prokaryotic or eukaryotic cell, in which the SOD gene can be expressed

and/or replicated. The host can then be cultured, after which any SOD, produced by it, can be recovered. Such a process for expressing and/or replicating an SOD gene can be carried out in a conventional manner, using known expression vectors and host environments (See Winnacker, 1987 and the references cited therein), as well as known replicons which can carry the SOD gene so that the gene is expressed, as well as propagated, in a host.

The Examples, which follow, illustrate the invention. In the Examples, reference is made to the accompanying drawings in which:

Figure 1: The aminoacid sequence (bottom) of the mature MnSOD of Nicotiana plumbaginifolia, linked to the mitochondrial transit peptide, and the corresponding DNA sequence coding for this amino acid sequence (top): This DNA sequence corresponds to the cDNA as comprised in plasmid pSOD-1. The numbering refers to the nucleotides of the open reading frame comprising the mitochondrial transit peptide and the mature MnSOD. The N-terminus of the MnSOD is indicated by an arrow.

Figure 2: The amino acid sequence (bottom) of the mature MnSOD of Nicotiana plumbaginifolia and the corresponding DNA sequence coding for this amino acid sequence (top) used in Example 4. The real N-terminus of the MnSOD is indicated by an arrow; the preceding amino acids are derived from the cloning procedure.

Figure 3: The amino acid sequence (bottom) of a polypeptide with MnSOD activity of Nicotiana plumbaginifolia, linked to the chloroplast transit peptide, and the corresponding DNA sequence coding for this amino acid sequence

19

(top) used in Example 5. The N-terminus of the mature MnSOD is indicated by an arrow. The expected cleavage site of the transit peptide is indicated by a double arrow.

Figure 4a: Restriction map of plasmid pDE9 containing the 35S3 promoter from CaMV isolate CabbbB-JI.

Figure 4b: DNA sequence of the 35S3 promoter fragment of CaMV isolate CabbbB-JI.

Figure 5: The amino acid sequence (bottom) of the mature FeSOD of Nicotiana plumbaginifolia, as cloned in the PstI site of plasmid PUC18, and the corresponding DNA sequence coding for this amino acid sequence (top) used in Example 12. The sequence is given for the lacZ initiation codon of pUC18. The start of the FeSOD sequence is indicated by an arrow.

Figure 6: Construction of pEX1SOD of Example 3.

Figure 7: Construction of pEX3SOD of Example 4.

Figure 8: Construction of pEX4SOD of Example 5.

Figure 9: Construction of pEX5SOD of Example 6.

Figure 10: Expression of MnSOD in transgenic plant calli.

Figure 11: Percent weight change in relation to paraquat concentration for leaf discs derived from transgenic (T16-213 (11A) and T16-202 (11B) overexpressing MnSOD in the chloroplasts) and control (T17-50) N. tabacum PBD6 (Crosses: transgenic plants; open circles: control plants).

Figure 12: Percent bleaching of chlorophyll pigments (measured at 664nm) in relation to paraquat concentration for leaf discs derived from transgenic (T16-213 (12A) and T16-202 (12B) overexpressing MnSOD in the chloroplasts) and control (T17-50) N. tabacum PBD6 (Crosses:

20

transgenic plants; open circles: control plants).

Figure 13: Percent bleaching of chlorophyll in relation to initial chlorophyll content (measured as E_{ini} per 35 mg of leaf tissue) for leaf discs derived from various transgenic and control N. tabacum PBD6 (treated with 50uM paraquat for 24 hrs.)

Figure 14: TBA reactivity (TBAR) in relation to paraquat concentration for various transgenic and control N. tabacum PBD6. Expression levels of MnSOD for each plant are indicated.

The abbreviations used in the Figures and in the Examples are:

Gluc:	Glucose
Sucr:	Sucrose
Mann:	Mannitol
mit-tp:	Mitochondrial transit peptide encoding sequence
cp-tp:	Chloroplast transit peptide encoding sequence
ori:	origin of replication (Ori-1: of pBR322 plasmid of <u>E. coli</u> , Ori-2: of pVS1 plasmid of <u>Pseudomonas aeruginosa</u>).
Sm:	Streptomycin resistance gene
Cont:	Control
MV:	Methyl Viologen (Paraquat)
Ch+, Ch++,	
Ch+++:	Expression levels of MnSOD in chloroplasts of transgenic plants
M+, M++,	
M+++:	Expression levels of MnSOD in mitochondria of transgenic plants
C+:	Expression levels of MnSOD in cytosol of

transgenic plants

Sp: Spectinomycine resistance gene
 RB-LB: Right and left borders of T-DNA of Agrobacterium tumefaciens
 5 35S: 35S promotor of CaMV isolate CM1841.
 35S3: 35S promotor of CaMV isolate CabbB-JI
 3'g7: polyadenylation signal sequence of T-DNA gene
 7
 pnos: promoter of the nopaline synthase gene of A. tumefaciens.
 10 3' ocs: polyadenylation signal sequence of the
 octopine synthase gene of A. tumefaciens.
neo: neomycine phosphotransferase II gene
 MnSOD: manganese superoxide dismutase gene of
 15 Nicotiana plumbaginifolia.
 cip: Calf intestinal alkaline phosphatase
 pSSU: promoter of the 1A small subunit of Rubisco of
Arabidopsis thaliana.

20 Unless otherwise specified in the Examples, all
 procedures for making and manipulating recombinant DNA are
 carried out by the standardized procedures described by
 Maniatis et al., 1982. The following plasmids and vectors,
 used or prepared in the Examples, have been deposited in
 25 the Deutsche Sammlung Für Mikroorganismen und Zellkulturen
 ("DSM"), Mascheroder Weg 1B, Braunschweig, Federal
 Republic of Germany under the provisions of the Budapest
 Treaty:

Plasmid or Vector	DSM Accession No.	Date of Deposit
pSC1701A2	DSM 4286	22 Oct. 1987
pGSC1700	DSM 4469	21 Mar. 1988
pEX1SOD	DSM 4695	8 Jul. 1988
pEX3SOD	DSM 4696	8 Jul. 1988
pEX4SOD	DSM 4692	8 Jul. 1988
pEX5SOD	DSM 4693	8 Jul. 1988

Example 1: Isolation of superoxide dismutase cDNA clone from *Nicotiana plumbaginifolia*.

Cell suspension cultures were initiated from homozygous *N. plumbaginifolia* plants. Total protein extracts were separated by two-dimensional polyacrylamide gel electrophoresis, and the separated proteins were recovered by electroblotting onto membranes that allow direct gas-phase sequencing analysis of the immobilized proteins (Bauw et al., 1987). Proteins were visualized by U.V. illumination after treatment with fluorescamine. Protein spots were removed by scissors and stored at -20°C (Bauw et al., 1987). One of the isolated proteins had the following NH₂- terminal sequence:

LQTFSLPDLPYDXGALEPAI?GD

in which "?" is an unknown residue and "X" indicates a modified residue (for standard abbreviations of amino acids see Singleton and Sainsbury, 1987). Comparison of the protein sequence with published eukaryotic protein sequences within the National Biomedical Research Foundation Protein Sequence Data Bank (Release 9) [U.S.A.] showed partial homology with the human Mn superoxide dismutase (Harris and Steinman, 1977) and with the corresponding enzyme of *Saccharomyces cerevisiae* (Harris and Steinman, 1977). The determined N-terminal sequence was, therefore, presumed to belong to a plant MnSOD. To obtain the cDNA clone encoding the complete enzyme, an oligonucleotide was designed to match part of the N-terminal amino acid sequence. This was synthesized with a deoxyinosine at ambiguous codon positions (Ohtsuka et al., 1985; Takahashi et al., 1985) and used as probe to screen a cDNA library from a *N. plumbaginifolia* cell

23

suspension culture depleted 14 days for cytokinin (Bauw et al., 1987). The procedure to construct the cDNA library was as described in Gubler and Hoffman (1983). The oligonucleotide sequence was:

5'-CCITAIGAITAIGGIGCTIGAIACCIGC-3'

5X10⁴ clones from the cDNA library were hybridized with the oligonucleotide at 40°C. Twelve clones showed a positive signal. One clone "pSOD1" was selected for further analysis. The pSOD1 cDNA was sequenced on both strands according to the method of Maxam and Gilbert (1980). The entire sequence of the 996 bp cDNA insert is shown Fig. 1, with its flanking G/C homopolymer tails added during the cloning procedure. It contains one continuous open-reading frame, corresponding to 228 amino acids. The sequence homologous to the oligonucleotide probe is underlined. The cDNA clone also contains a mitochondrial leader sequence of 24 amino acids upstream from the mature protein (Fig. 1). The amino acid sequence, deduced from the cDNA sequence, is written in Fig. 1 below the nucleotide sequence in the one-letter code. The amino acid sequence starting from amino acid 25 (indicated by arrow) is completely homologous to the previously determined N-terminal amino acid sequence of the mature protein. The molecular weight calculated from the cloned sequence is 22.8 kD for the mature MnSOD and 25.5 kD for the transit peptide-MnSOD preprotein.

A comparison between the *N. plumbaginifolia* mature protein and the MnSOD of bacteria, yeast and humans shows considerable homology (Bowler et al., 1989a).

The comparative homologies show that the plant MnSOD is more closely related to human and yeast, than to bacterial, MnSOD. However, a posteriori comparative

analysis of the respective genes shows that isolation of the plant MnSOD would have been difficult using a cloned bacterial, yeast or human SOD gene fragment as a probe for screening a plant cDNA library.

5 The presence in the signal sequence of the five arginine residues distributed among uncharged amino acids, the absence of acidic residues, and the occurrence of hydroxylated amino acids such as serine and threonine are typical for a leader sequence for translocation to the
10 mitochondrial matrix (Schatz, 1987). A mitochondrial location is consistent with data from analysis of subcellular fractions.

15 Example 2: Expression of MnSOD in *N. plumbaginifolia* plants.

MnSOD was identified as a highly abundant protein in cell suspension cultures of *N. plumbaginifolia*. Northern analysis on total RNA, isolated from different tissues of
20 *N. plumbaginifolia* plants, with the pSOD1 cDNA as probe revealed great variations in steady state mRNA. Plants were sterile-grown at 25°C on Murashige and Skoog ("MS") medium (Murashige and Skoog, 1962). They were grown with a 16-hour light/ 8-hour dark cycle on solid MS medium containing 0.1 M sucrose. Cell suspension cultures were
25 grown in the dark, in liquid medium with 0.1 M sucrose and supplemented with 0.5 mg/l naphthalene acetic acid ("NAA") and 0.1 mg/l 6-benzylaminopurine ("BAP"). They were subcultured every 3 days. Total RNA was prepared according to Jones et al. (1985). 12 ug RNA was denatured in
30 formaldehyde, electrophoresed, transferred to nylon membranes and hybridized with ³²p-labeled RNA-probes corresponding to the HpaI-Hind III fragment of pSOD1. Expression was very weak in leaves of intact plants, 2- to 3-fold higher in roots, and 50 times higher in dark-grown
35

cell suspension cultures (Bowler et al., 1989a). These differences in expression were not due to a switch-off of photosynthesis, since exposure of whole plants to the dark did not result in increased expression of MnSOD. Treatment with several concentrations of paraquat resulted in very minor increases of MnSOD mRNA in leaves (Bowler et al., 1989a). Incubation of leaf discs for 48 hours in the dark in liquid medium yielded induced MnSOD mRNA levels comparable to those in cell suspension cultures. Incubation of leaf discs in pure water caused only a 2-3 fold induction, which is similar to the effect of wounding whole plants by cutting their leaves in several places (Bowler et al., 1989a).

Sucrose was found to be a crucial factor for induction of MnSOD RNA. The effect was greatest in the presence of salts and showed a linear dose-response at sucrose concentrations ranging from 0.001 till 0.2 M. The highest level of expression was reached after 48 hrs incubation. Combinations of salts delivering iron, manganese, copper and zinc ions in the presence of sucrose showed that the induction was not due to these salts in particular. The increase in MnSOD mRNA was also possible by induction with glucose, but not with mannitol (Bowler et al., 1989a).

Levels of MnSOD protein were measured by assaying SOD activity. Leaf discs were incubated for various time periods in MS + 0.1 M sucrose supplemented with 0.5 mg/l NAA and 0.1 mg/l BAP and subsequently homogenized in an equal volume of cold extraction buffer (50 mM potassium phosphate, pH 7.8, 0.1 % ascorbate, 0.05 %, beta-mercaptoethanol, 0.2 % Triton X-100) and centrifuged at 13,000 rpm for 12 minutes. Protein samples were separated on non-denaturing 10 % polyacrylamide gels run at 120 V constant voltage. SOD activity was localized on these gels using the in situ staining technique according

to Beauchamp and Fridovich (1971). Inhibition studies with H_2O_2 and KCN (Bridges and Salin, 1981) revealed the upper band to be MnSOD and the lower band (only present in samples containing more than 100ug protein) a Cu/ZnSOD. Semi-quantitative data were obtained by loading several concentrations (10-200 ug) of total protein on the gel. Change of MnSOD activity was best visualized on samples containing 50 ug protein. In relation to the mRNA profile, a similar, albeit delayed, induction pattern was obtained at the protein level (Bowler et al., 1989a). The increase in SOD activity caused by addition of exogenous sugars appeared to be entirely due to expression of MnSOD, since the Cu/ZnSOD in the extracts showed no significant alteration in expression level (Bowler et al., 1989a).

Example 3: Construction of pEX1SOD comprising the MnSOD cDNA under the control of the CaMV 35S promoter (Fig. 6)

The MnSOD cDNA was isolated from pSOD1 of Example 1 as a 910 bp HpaI-SmaI restriction fragment. This fragment was cloned in the vector PGSJ780A derived from pGSC1701A2. Plasmid PGSJ780A contains the 35S promoter fragment (Odell et al., 1983) from CaMV isolate CM1841 (Gardner et al., 1981) and the 3' untranslated region of T-DNA gene 7 (Velten and Schell, 1985) as well as a chimaeric cassette for plant transformation selection. This vector was digested with ClaI and treated with Klenow DNA polymerase to generate blunt ends. This yielded the plasmid "pEX1SOD" which contains, between T-DNA border sequences, two chimaeric genes:

The first chimaeric gene contains:

- the CaMV 35S promoter,
- the MnSOD gene consisting of the coding sequences for both the mitochondrial transit peptide and the mature MnSOD (Figure 1), and

27

- the 3' end of T-DNA gene 7 used for correct polyadenylation of the mRNA;

The second chimaeric gene contains:

- the nopaline synthase promoter,
 - the neo gene encoding neomycin phosphotransferase II, and
 - the 3' end of octopine synthase,
- and serves as a selectable marker during plant transformation (Hain et al., 1985). Since the plasmid contains a plant MnSOD gene with its own transit peptide sequence under the control of the constitutively expressed 35S promotor, targeting to the mitochondria is expected.

Example 4: Construction of pEX3SOD comprising the MnSOD cDNA sequence lacking the mitochondrial signal peptide sequence under control of the CaMV 35S promoter (Fig. 7)

The SacII site, which is present at the transit peptide cleavage site of the MnSOD clone in pSOD1 from Example 1, was converted to a BglII site as follows. pSOD1 was digested with SacII, blunt ended with Klenow DNA polymerase and ligated with octamer BglII linkers. This yielded the plasmid "pSOD-B". The BglII-BamHI fragment from pSOD-B, containing the SOD cDNA clone, was isolated and cloned in the BamHI site of pGSJ780A. This yielded the plasmid "pEX3SOD" which contains, between T-DNA border sequences, two chimaeric genes:

The first chimaeric gene contains:

- the CaMV 35S promoter,
- the MnSOD encoding sequence without transit peptide encoding sequence (Figure 2), and
- the 3' end of gene 7.

The second chimaeric gene contains:

- the nopaline synthase promoter,
- the neo gene, and

- the 3' end of octopine synthase,
and serves as a selectable marker for plant
transformation. The MnSOD cDNA, under the control of the
35S promoter, will yield a cytoplasm localized MnSOD,
constitutively expressed in all tissues.

Example 5: Construction of pEX4SOD comprising the MnSOD
cDNA sequence and an upstream chloroplast transit peptide
sequence under control of the CaMV 35S promoter (Fig. 8)

The BglIII-BamHI fragment was isolated from pSOD-B of
Example 4 and cloned in the BamHI site of pKAH1 yielding
pKAH1-SOD. pKAH1 contains the transit peptide sequence of
the Rubisco pea small subunit ("ssu") gene SS3.6
(Cashmore, 1983). pKAH1-SOD contains the SOD cDNA clone
(without mitochondrial transit peptide) fused to the
chloroplast transit peptide ("tp") sequence of the ssu
gene under the control of the 35S promoter of CaMV isolate
CM1841. Finally, the BglIII-BamHI fragment from pKAH1-SOD,
carrying the 35S-tp-SOD cassette, was cloned in the BamHI
site of pGSC1702 yielding the plasmid "pEX4SOD". pGSC1702
is derived from pGSC1700 and contains the 3' untranslated
region of T-DNA gene 7 as well as a chimaeric cassette for
transformation selection. The chimaeric construct in
pEX4SOD differs from that in pEX3SOD in having a
chloroplast transit peptide sequence at the N-terminus of
the MnSOD cDNA coding sequence as shown in Figure 3. In
this regard, the mitochondrial transit peptide sequence
has been replaced by a chloroplast transit peptide
sequence in order to constitutively express a MnSOD which
is targeted to chloroplasts.

Example 6: Construction of pEX5SOD comprising the MnSOD cDNA under the control of the Arabidopsis Rubisco small subunit promoter (Fig. 9)

5 The BglIII-EcoRI fragment of pSOD-B, containing the SOD fragment, was cloned into the plasmid pC23, digested with BglIII and EcoRI, yielding the plasmid "pC23SOD-B". pC23 was obtained from pC22 (Simoens et al., 1986) by the downstream extension of the polylinker

10 ApaI-SmaI-EcoRI-XbaI by a StuI, a HindIII and a BglIII site. As described in European patent 242,246, a Rubisco small subunit gene ("ssu") promoter from Arabidopsis thaliana is contained in plasmid pATS3 as a 1.5 kb EcoRI-SphI fragment. The ssu promoter fragment was cloned

15 into a vector (designated as pGS1400) so that it could be excised as a BglIII-BamHI fragment. This BglIII-BamHI fragment was cloned in the BglIII site of pC23SOD-B, yielding the plasmid "pC23SSUSOD". Finally, the BglIII-BamHI fragment from pC23SSUSOD, containing the ssu promoter fragment fused to the SOD cDNA clone, was

20 isolated as a BglIII-BamHI fragment and inserted into the BamHI site from pGSC1702 upstream of the 3' end of gene 7 yielding the plasmid "pEX5SOD". This plasmid contains, between T-DNA border sequences, two chimaeric constructs:

One chimaeric gene contains:

- 25
- the Rubisco small subunit 1A promoter from Arabidopsis thaliana,
 - the MnSOD mature protein coding sequence, and
 - the 3' end of gene 7.

The second chimaeric gene contains:

- 30
- the nopaline synthase promoter,
 - the neo gene, and
 - the 3' end of octopine synthase,

and serves as a selectable marker during plant transformation. Since the leaderless MnSOD cDNA sequence

35

is under the control of a ssu gene promoter, light regulated expression of the MnSOD in the cytoplasm is expected.

5 Example 7: Chimaeric constructs under the control of the 35S3 promoter from the CaMV isolate CabbB-JI

The 35S3 promoter of CaMV isolate CabbB-JI (Hull and Howell, 1978) was cloned in pUC19 (Yannish-Perron et al., 1985) yielding pDE9 (Fig. 4A) and sequenced. The sequence
10 of the 35S3 promoter fragment as contained in pDE9 is presented in Fig. 4B. The NcoI site of the promoter fragment was created at the first ATG codon occurring in the 35S3 RNA transcript by site directed mutagenesis using the pMa5-8 and pMc5-8 plasmids (European patent
15 application 87402348.4) and the gapped duplex procedure of Stanssens et al. (1987).

Analysis of the nucleotide sequence of the 35S3 promoter showed that it differs from that of the 35S promoter from CaMV isolate CabbB-S (Franck et al., 1980)
20 and from CaMV isolate CM1841 (Gardner et al., 1981). Moreover, some chimaeric constructs with the 35S3 promoter have shown greater activity in transgenic plants than with the 35S promoter from CabbB-S (Harpster et al., 1988). The 35S3 promoter can be clearly distinguished from the other
25 two 35S promoters, mentioned above, by the absence of an EcoRV site due to a single nucleotide substitution, immediately ahead of the TATA box.

Using standard recombinant DNA techniques, the 35S promoter in the plant expression vectors pEX1SOD, pEX3SOD
30 and pEX4SOD of Examples 3, 4 and 5 is replaced by the 35S3 promoter to yield three additional plant expression vectors "pEX6SOD", "pEX7SOD" and "pEX8SOD", respectively.

Example 8: Plant transformation and regeneration

Using well-known techniques as described in European patent publication 116718 and European patent application 87400544.0, plants cells are transformed, as described below, with the plant expression vectors pEX1SOD, pEX3SOD, pEX4SOD, pEX5SOD, pEX6SOD, pEX7SOD and pEX8SOD from Examples 3-6 and 7.

The plant expression vectors are mobilized into the Agrobacterium tumefaciens recipient strain C58C1Rif (pGV2260) according to the procedure described by Deblaere et al. (1985). These strains are then used for the transformation and regeneration of: Arabidopsis thaliana according to the procedure of Valvekens et al., (1988); and Nicotiana plumbaginifolia and Nicotiana tabacum PBD6 according to the leaf disc transformation procedure of De Block et al. (1987). Transformed shoots of N. plumbaginifolia and Nicotiana tabacum are regenerated into whole plants according to the methods of Ellis et al. (1988) and De Block et al. (1987), respectively.

Example 9: Analysis of transgenic plants

The plants of Example 8, transformed with pEX1SOD, pEX3SOD and pEX4SOD, were analyzed for the expression of recombinant SOD genes as follows.

Transgenic Nicotiana plumbaginifolia calli were homogenized, and MnSOD activity was assayed by in situ staining after electrophoresis in non-denaturing polyacrylamide gels, using homogenization and electrophoresis procedures as described in Example 2. The gels were treated with KCN and H₂O₂ prior to staining so that only the manganese isoforms of the enzyme were revealed. The results of these assays are presented in Figure 10 in which the MnSOD activity in control calli and calli transformed with pEX1SOD, pEX3SOD and pEX4SOD can be seen.

5 The endogenous MnSOD is clearly marked. The lane for pEX1SOD shows the 35S-mit-tp-MnSOD construction to be expressed and to be targeted to the mitochondria as the additional MnSOD comigrates with the endogenous enzyme. This results in a band of approximately double intensity when compared to the control.

10 The lane for pEX3SOD shows a faint band below the endogenous band. This band represents the MnSOD as expressed in the cytoplasm.

15 The lane for pEX4SOD shows two additional bands, one of which is at the same position as the faint band in the pEX3SOD lane. This represents the MnSOD expressed in the chloroplasts which suggests that the SSU transit peptide is cleaved off as expected. The other faint band in this lane, with an intermediate position with respect to the endogenous band and the processed form described above, may represent an unprocessed form.

20 The results of these MnSOD activity assays in transgenic calli prove that the recombinant SOD genes, which were introduced, were actively expressed. The expression level of both pEX1SOD and pEX4SOD is equivalent to that of the endogenous SOD gene as expressed in calli while the expression level of pEX3SOD is approximately one-tenth of that of the endogenous SOD gene. As the
25 endogenous MnSOD is highly expressed in calli, the chimaeric constructions give similarly high expression.

30 Similar results were obtained for leaf tissue of transformed *N. tabacum* PBD6. Moreover, in PBD6 plants, the expression levels as determined by SOD activity staining were found to correspond well with mRNA levels determined by Northern blotting using the cDNA of the MnSOD as a probe and with protein levels determined either by sodium dodecyl sulphate ("SDS") polyacrylamide gel electrophoresis (Laemmli, 1970) or by Western blotting
35

using polyclonal antisera raised against the MnSOD overexpressed in yeast (See Bowler et al., 1989b for expression of MnSOD in yeast).

To see whether the MnSOD expression was efficiently targeted to the desired cell compartments, leaf tissue of plants transformed with pEX1SOD, pEX3SOD and pEX4SOD was homogenized, and subcellular fractions were prepared. Fractions representing the chloroplast stroma and membranes were prepared according to Van den Broeck et al. (1985) and Mullet and Chua (1983), while fractions corresponding to the mitochondrial matrix and membranes were prepared according to Boutty et al. (1987). In addition, in situ immunolocalization of the MnSOD was performed on thin sections prepared from leaf tissue of transgenic plants (Greenwood and Chrispeels, 1985), using polyclonal antisera raised against the MnSOD (prepared as described above).

Both approaches showed that MnSOD activity in PBD6 plants transformed with pEX4SOD was exclusively localized in the chloroplasts, as expected. Total SOD activity in the chloroplasts of these plants was approximately doubled.

Mitochondrial targeting was also shown to be very effective in PBD6 tobacco plants transformed with pEX1SOD since MnSOD activity was exclusively localized within the mitochondria. Total SOD activity in these organelles was increased 20-fold with respect to nontransformed control plants.

Cytosolic expression was shown to be the least effective. Although total SOD activity in the cytosol of PBD6 plants transformed with pEX3SOD was not significantly increased as compared to control plants, the presence of MnSOD activity could be unequivocally determined while no increased MnSOD activity was found in the chloroplasts and

mitochondria.

Example 10: Tolerance of transgenic plants to increased levels of superoxide radicals.

5 In order to observe the tolerance of the transgenic plants of Example 9 to increased levels of superoxide radicals, plants were subjected to various concentrations of methyl viologen ("MV"). This substance is known to generate superoxide radicals, especially in chloroplasts, and its effects have been well characterized. In general, superoxide radicals, generated by the action of MV, peroxidize the lipids in biological membranes, initiating a chain propagation reaction. In chloroplasts, this eventually leads to an oxidation of the photosynthetic pigments in the membranes (Halliwell, 1984). Experiments with MV therefore provide a convenient model system to study alterations in the physiological state of transgenic plants overexpressing the MnSOD. Results are expected to have a direct bearing on situations in which plants are subjected to natural, superoxide radical-producing, stress conditions.

15 In general, leaf pieces of transformed and untransformed plants were incubated in Petri dishes containing various amounts of MV. To stimulate electron transport, carbonylcyanide p-trifluoromethoxyphenylhydrazine ("FCCP") was also included. After incubation, factors such as weight changes (due to membrane damage), pigment bleaching and the extent of lipid peroxidation were assayed and compared.

30 1. Weight Changes

Leaf discs of constant size (7x22 mm) and identical age were incubated in the light in aqueous solutions containing 2uM FCCP and various concentrations of MV for

35

21 hours at 20°C. The leaf discs were weighed before and after incubation. Percent weight change for two transgenic plants (T16-213 and T16-202) as compared to a control plant (T17-50) for different concentrations of MV is shown in Fig. 11. Each measurement was made for two separate leaf discs of the same plant and curves were drawn between the averages of the paired values.

The results clearly show the protective effect of chloroplastic MnSOD overexpression against loss of membrane integrity due to the action of MV.

2. Chlorophyll bleaching.

Incubation of leaf discs was performed as described above. After incubation, a pigment extract of the leaf discs was prepared as follows. Discs were homogenized in a mortar and pestle using Al_2O_3 as an abrasive. After addition of 3.1 ml of a chloroform/methanol/water mixture (1:2:0.1) and mixing, the suspension was allowed to stand in the dark for 5 minutes. The supernatant was transferred to a centrifuge tube, and 2.4 ml of a chloroform/methanol mixture (1:2) was added to the mortar. After a new extraction period of 5 minutes in the dark, the supernatant in the mortar was added to the centrifuge tube. The combined supernatants were then centrifuged for 15 minutes at 3300 rpm (Sorvall HB4 rotor) at room temperature (25°C). The extinction of the supernatant was measured at 664 nm (E_{final}). E_{664} was also determined for pigment extracts of comparable leaf discs without incubation (E_{ini}). Fig. 12 shows "percent bleaching" $[\%E = 100 \times (E_{\text{ini}} - E_{\text{fin}}) / E_{\text{ini}}]$ as a function of the MV concentration.

To eliminate possible effects of differences in initial chlorophyll content of the discs derived from different plants, a second experiment was performed as

follows. Leaf discs from different plants were incubated for one day in an aqueous solution containing 50 μ M MV. Percentage bleaching was calculated and standardized with respect to the initial amounts of chlorophyll present in each plant. The results are shown in Fig. 13. It can be seen that %E decreases linearly with increasing initial chlorophyll content (as measured by E_{ini} per 35 mg of chlorophyll) at least for control (T17-50 and PBD6), cytosolic MnSOD (T16-100 and T16-109) and mitochondrial MnSOD plants (T16-7 and T16-37). Chloroplastic MnSOD plants (T16-202 and T16-213), by comparison, are well below this line, indicating a protection against pigment oxidation in these plants.

3. Lipid peroxidation.

The assay for the extent of lipid peroxidation of polyunsaturated fatty acids is based on the detection of malondialdehyde, a decomposition product of lipid peroxides. Production of malondialdehyde is enhanced by acidic conditions. Malondialdehyde reacts with thiobarbituric acid ("TBA") to produce a red chromogen which can be measured photometrically at 532 nm (Slater, 1984). The increase in TBA reactivity ("TBAR") is thus a direct measure of lipid peroxidation.

PBD6 leaf discs (0.4 g) of approximately 1 cm² were incubated in the light at room temperature in petri dishes containing 10 ml 50 mM Tris/HCl, pH 7.0, with different concentrations of MV (0, 0.1, 0.5 and 1.0 mM). After a two hour incubation period, tissue was homogenized in: 3.5 ml 6 mM NaH₂PO₄; 1.2 mg/l ethylenediamine tetraacetate ("EDTA"); and 0.265% (w/v) TBA in 0.17 M HCl. The homogenate was placed in a boiling water bath for 15 minutes, cooled to room temperature and centrifuged at 12000 rpm for 10 minutes (Eppendorf centrifuge). The

absorbance of the supernatant was read at 532 nm and converted to the amount of malondialdehyde per gram of fresh tissue.

TBAR for all transgenic plants was less than for control PBD6 plants (Figure 14A). Chloroplastic MnSOD plants are clearly best protected against lipid peroxidation, and this effect could be shown to be dependent upon the expression level of the MnSOD (see Fig. 14B).

Example 11: Tolerance of transgenic plants to stress conditions.

The response of the transgenic plants of Example 9 to different stress situations was analyzed in depth through monitoring of fitness components such as growth and survival characteristics. Comparison is made to suitable controls (i.e., non-transformed plants). Environmental differences and/or differences in genetic background among the tested plants necessitate a statistical approach.

In one case, the germination of seeds of Arabidopsis plants transformed with pEX1SOD, PEX3SOD and PEX4SOD was assayed under conditions of constant light. Seeds of transformed and untransformed plants were germinated on K1 medium (Valvekens et al., 1988) in petri dishes sealed with two layers of Urgopore^(R) porous tape (from Chenove Co. in France) and placed under conditions where the light intensity was held constant at 2500 lux. The temperature of the growth chamber was set at 21°C. The experiment was initiated on January 26, 1989 (winter), and due to the configuration of the growth chamber, chamber temperature was somewhat influenced by outside temperatures and fluctuated between 10°C and 20°C in phase with the outside temperatures.

For each tested Arabidopsis line, three K1 plates,

with 100 seeds each, were placed in the chamber. Plates were randomly spaced in sets of three. After 15 days, the condition of the seedlings was assayed as follows: 0, not germinated; 1, germinated but seedlings show obvious signs of bleaching and/or retardation (indicative of environmental stress); and 2, germinated to normal seedlings. Further analyses were only performed for germinated seeds. Table 1 below displays, for each plant, the total number of seedlings scored in the two classes. For each plant, the value of chi square ("X²" -- Snedecor and Cochran, 1980) was calculated with respect to the results from the control plant (nontransformed Arabidopsis). From Table 1, it can be seen that all transgenic plants (i.e., 1SOD-, 3SOD- and 4SOD-plants transformed with pEX1SOD, pEX3SOD and pEX4SOD, respectively) have more normal seedlings (class 2) than the control plant. In all cases, the differences are highly significant.

Table 1

Arabidopsis line	Class 1 seedlings	Class 2 seedlings	X ²
C24 (control)	151	18	
1SOD4	54	71	72.5
1SOD5	52	175	194.7
1SOD7	38	204	241.9
3SOD5	106	49	37.3
3SOD11	103	76	80.5
4SOD1	58	186	252.6
4SOD7	65	124	112.6
4SOD2	65	122	110.9

Surface sterilized seeds of Arabidopsis thaliana (both untransformed and transformed with pEX1SOD, pEX3SOD or pEX4SOD) are germinated on germination medium as described by Valvekens et al. (1988). In the case of transgenic plants, the medium is supplemented with 50 mg/l kanamycin sulphate. One week old seedlings are transferred to germination medium supplemented with Fe(II)-EDTA at concentrations of 500 μ M, 250 μ M and 100 μ M. The plants are grown according to Valvekens et al. (1988), and the tolerance of transformed plants to the toxic effects of Fe (II) appears to be better than the tolerance of untransformed (wild-type) plants.

Example 12: Isolation of a cDNA clone from Nicotiana plumbaginifolia encoding an FeSOD.

cDNAs prepared from mRNAs, derived from a suspension culture of Nicotiana plumbaginifolia (see Example 1), were cloned in the PstI site of plasmid pUC18 (Yannisch-Perron et al., 1985) by means of homopolymeric dG-dC tailing as follows. pUC18 was linearized with PstI and a dG tail was added to the 3' ends by means of terminal deoxynucleotidyl transferase. Complementary dC-tails were added to the 3' ends of the cDNAs. 1 μ g of the resulting cDNA library was electroporated in 10^9 cells of an E.coli strain sodA⁻sodB⁻ (Carlioz and Touati, 1986) by means of the Biorad Gene Pulser (BioRad Chemical Division, 1414 Harbour Way, South Richmond, California 94804, U.S.A.) according to the procedure described in the BioRad manual. E.coli sodA⁻sodB⁻ is deficient in SOD activity and is unable to grow aerobically on minimal medium. Consequently, cDNA clones encoding SOD enzymes can be isolated by growing transformants on minimal medium. After 3 days of incubation at 37°C, a total of 67 colonies could be picked up. The inserts of some of the clones did not hybridize

with the cDNA coding for the MnSOD of Example 1. Restriction analysis of these clones showed that many contained very similar inserts. Proteins were extracted from colonies containing these plasmids, and assays for SOD activity on polyacrylamide gels (Beauchamp and Fridovich, 1971) confirmed that these colonies synthesized a protein with SOD activity. Inhibition studies (Bowler et al., 1989 a) further showed the SOD to be resistant to KCN and sensitive to H_2O_2 , results which are indicative of an FeSOD (Bannister et al., 1987). The pUC18 insert was sequenced (Sanger et al., 1977), and the DNA sequence is shown in Fig. 5. This is the first cDNA of an FeSOD to be isolated from an eucaryote. The first in frame codon after the dC-dG tail is a lysine, indicating that transcription is initiated from the lacZ-ATG codon of pUC18 and that the FeSOD is synthesized as a fusion protein. The FeSOD in leaf tissue of Nicotiana plumbaginifolia was found to be localized within the chloroplasts and to be highly responsive to stress conditions like MV, heat shock, sucrose, and infection by Pseudomonas syringae.

Plant expression vectors containing the FeSOD gene are prepared using the procedure of Examples 3-6, and such vectors are used to transform the same plant species as in Example 8, using the general procedures of Example 8, whereby the plants express the FeSOD gene.

Needless to say, this invention is not limited to the transformation of any specific plant(s). The invention relates to any plant, the genome of which can be transformed with an SOD gene, particularly a plant MnSOD or FeSOD gene, that is under the control of a promoter capable of directing expression of the SOD gene in the plant's cells and that is preferably fused at its 5' end to a targeting sequence encoding a targeting peptide for translocation within, or secretion from, the cells of the

41

plant of an expressed SOD, to provide the plant with increased resistance and/or tolerance to a naturally occurring stress condition which produces toxic, highly reactive, oxygen species in one or more of the plant cell compartments.

This invention also is not limited to the plant SOD genes of Figures 1, 2, 3 and 5, used in the foregoing Examples. In this regard, the invention encompasses MnSOD and FeSOD genes encoded by: 1) any DNA fragments differing from the SOD genes of Figures 1, 2, 3 and 5 by the replacement of any of their nucleotides by others, without modifying their genetic information (normally within the meaning of the universal genetic code); and 2) any DNA fragments that encode polypeptides which have the same or equivalent SOD properties as the polypeptides, encoded by the SOD genes of Figures 1, 2, 3 and 5, but which may not have the same amino acid residues. Likewise, this invention is not limited to the MnSODs and FeSODs of Figures 1, 2, 3 and 5 but rather covers any equivalent polypeptides. Indeed, it is apparent that one skilled in the art will be able to remove 5' and/or 3' portions of the SOD genes of Figure 1, 2, 3 and 5 without significantly affecting their usefulness for transforming plants to render them stress resistant in accordance with this invention. Such portions may be removed, for example, by removing terminal parts on either side of a SOD gene with an exonucleolytic enzyme (e.g., Bal31), and the remaining shortened DNA fragment can then be recovered in a suitable plasmid so that the capacity of the modified plasmid to transform plant cells and to enhance SOD production therein (e.g., as measured by the assay described in Example 2) can be determined. Such a shortened DNA fragment, coding for a shortened SOD which retains its SOD activity, is considered an equivalent of

42

an SOD gene of Figure 1, 2, 3 or 5. Likewise, such a shortened SOD is considered an equivalent of an SOD of Figure 1, 2, 3 or 5.

5 Furthermore, this invention is not limited to the promoters, 3' transcription regulation signals and targeting sequences used in the Examples. One skilled in the art will be able readily to substitute different DNA fragments and regulatory sequences which can perform
10 equivalent functions in the recombinant SOD gene of this invention in a transformed plant cell, cell culture or plant.

Also, this invention is not limited to the specific plasmids and vectors described in the foregoing Examples, but rather encompasses any plasmids and vectors containing
15 the recombinant SOD gene of this invention, useful for obtaining expression of the SOD gene in one or more plant cell compartments.

20

25

30

35

References

- ARRIGONI, ZACHEO, BLEVE-ZACHEO, ARRIGONI-LISO and LAMBERTI, 1981, Nematol. Medit. 9:195-198
- 5 - ASADA, KISO and YOSHIKAWA, 1974, J. Biol. Chem. 247:2175-2181
- BANNISTER, BANNISTER and ROTILIC, 1987, CRC Crit. Rev. Biochem. 22:111-180
- 10 - BAUW, DE LOOSE, INZE, VAN MONTAGU and VANDEKERCKHOVE, 1987, PNAS USA* 34:4806-4810
- BARRA, SCHININA, SIMMACO, BANNISTER, BANNISTER, ROTILIO and BOSSA, 1984, J. Biol. Chem. 259:12595-12601
- BEAUCHAMP and FRIDOVICH, 1971, Anal. Biochem. 44:276-287
- 15 - BEEVERS, 1979, Ann. Rev. Plant Physiol. 30:159-193
- BIRMINGHAM-McDONOGH, GRALLA and VALENTINE, 1988, PNAS USA 85:4789-4793.
- BOUTRY, NAGY, POULSEN, AOYAGI and CHUA, 1987, Nature 328:340-342
- 20 - BOWLER, ALLIOTTE, DE LOOSE, VAN MONTAGU and INZE, 1989a, EMBO J. 8:31-38
- BOWLER, ALLIOTTE, VAN DEN BULCKE, AUW, VANDEKERCKHOVE, VAN MONTAGU and INZE, 1989b, PNAS USA 3237-3241
- BRIDGES and SALIN, 1981, Plant Physiol. 68:275-278
- 25 - BROCK and WALKER, 1980, Biochemistry 19:2873-2882
- BROWN and FRIDOVICH, 1981, Arch. Biochem. Biophys. 206:414-419

* Proceedings of the National Academy of Sciences, USA

- CANNON, WHITE and SCANDALIOS, 1987, PNAS USA 84: 179-183
- CARLIOZ LUDWIG, STALLINGS, FEE, STEINMAN, TOUATI, 1988, J. Biol. Chem. 263:1555-1562
- 5 - CARLIOZ and TOUATI, 1986, EMBO J. 5:623-630
- CASHMORE, 1983, in "Genetic Engineering of Plants" (Kosugen, Meredith and Hollaender, eds.) Plenum Press.
- CLARE, RABINOWITCH and FRIDOVICH, 1984, Arch. Biochem. Biophys. 231:158-163
- 10 - COLLINGE and SLUSARENKO, 1987, Plant. Mol. Biol. 9:389-410
- DAUB, 1982, Plant Physiol. 69:1361-1364
- DAUB, 1987, Phytopathology 77:1515-1520
- DAUB and BRIGGS, 1983, Plant Physiol. 71:763-766
- 15 - DAUB and HANGARTER, 1983, Plant Physiol. 73:855-857
- DEBLAERE et al., 1985, NAR* 13:4777-4788
- DEBLOCK et al., 1987, EMBO J. 6:2513-2518
- ELLIS et al., 1988, Plant Molecular Biol. 10:203-214
- 20 - FRANCK, GUILLEY, JONARD, RICHARDS and HIRTH, 1980, Cell 21:285-294
- FRIDOVICH, 1987, Science 201:875-880
- FUCCI, OLIVER, COON and STADTMAN, 1983, PNAS USA 80:1521-1525
- 25 - FURUSAWA and MIZUGUCHI, 1988, 5th Intl. Cong. of Plant Pathology, Aug. 20-24, 1988, Kyoto, Japan, Abstracts of Papers 2.1.
- FURASAWA, TANAKA, THANUTONG, MIZUGUCHI, YAZAKI and ASADA, 1984, Plant Cell Physiol. 25:1247-1254
- 30 - GARDNER, HOWARTH, HAHN, BROWN-LUEDI, SHEPARD and MESSING, 1981, NAR 9:2871-2887

*Nucleic Acids Research

- GETZOFF, TAINER, STEMPIEN, BELL and HALLEWELL, 1989, Proteins, 5:322-336
- GIELEN, DE BEUCKELEER, SEURINCK, DEBOECK, DE GREVE, LEMMERS, VAN MONTAGU and SCHELL, 1984, EMBO J. 3:835-845
- GREENWOOD and CHRISPEELS (1985) Plant Physiol. 79: 65-71
- GRIMES, PERKINS and BOSS, 1983, Plant Physiol. 72:1016-1020
- GUBLER and HOFFMAN, 1983, Gene 25:263-269.
- HAIN, STABEL, CZERNILOVSKY, STEINBISS, HERRERA-ESTRELLA and SCHELL, 1985, Mol. Gen. Genet. 199:161-168
- HALLIWELL, 1984, "Chloroplast Metabolism", Clarendon Press, Oxford
- HARBOUR and BOLTON, 1975, Biochem. Biophys. Res. Commun. 64:803-807
- HARPSTER, TOWNSEND, JONES, BEDBROOK and DUNSMUIR, 1988, Mol. Gen. Genet. 212:182-190
- HARRIS and STEINMAN, 1977, in "Superoxide and superoxide dismutase", Michelsen, McCord and Fridovich (eds.), Academic Press, London, p225-230
- HARTMAN, SUZUKI and STACK, 1989, Appl. Environmental Microbiol. 1989:7-14
- HASSAN and FRIDOVICH, 1979, Arch. Biochem. Biophys. 196:385-395
- HO and CRAPO, 1988, FEBS 229:256-260
- HOHN and SCHELL, 1987 (eds.) Plant Infectious Agents. Springer Verlag Wien, New York.
- HULL and HOWELL, 1978, Virology 86:482-493
- IMLAY and LINN, 1988, Science 240:1302-1309
- JONES, DUNSMUIR and BEDBROOK, 1985, EMBO J. 4:2411-2418
- KAISER, 1979, Planta 145:377-382
- KUHLEMEIER, GREEN and CHUA, 1987, Ann. Rev. Plant. Physiol. 38:221-257

- LAEMMLI, 1970, Nature, 227:680-685
- LEE and BENNETT, 1982, Plant Physiol. 69:1444-1449
- LEE, BOCHNER and AMES, 1983, PNAS USA 80:7496-7500
- 5 - MACKAY, SENARATNA, MCKERSIE and FLETCHER, 1987, Plant
Cell Physiol. 28:1271-1278
- MANIATIS, FRITSCH and SAMBROOK, 1982, Molecular Cloning
- A laboratory Manual, Cold Spring Harbor Laboratory,
Cold Spring Harbor
- 10 - MARRES, VAN LOON, OUDSHOORN, VAN STEEG, GRIVELL and
SLATER, 1985, Eur. J. Biochem. 147:153-161.
- MATTERS and SCANDALIOS, 1986, Biochem. Biophys. Acta
882:29-38
- MAXAM and GILBERT, 1980, Methods Enzymol.
15 65(1):499-559, Academic Press
- MEAD, 1976, in "Free Radicals in Biology Vol 1" (Pryor,
ed) pp. 51-68 Academic Press
- MONK, FAGERSTEDT and CRAWFORD, 1987, Plant Physiol.
85:1016-1020
- 20 - MULLET and CHUA, 1983, Meth.Enzymol. 97:502-509
- MUNKRES, RANA and GOLDSTEIN, 1984, Mechanisms of Ageing
and Development 24:83-100
- MURASHIGE and SKOOG, 1962, Physiol. Plantarum
15:473-497
- 25 - NAGY and MALIGA, 1976, Z. Pflanzenphysiol. 78:453-455
- ODELL et al., 1983, Nature 313:810-812
- OHTSUKA, MATSUKI, IKEHARA, TAKAHSHI, MATSUBARA, 1985,
J. Biol. Chem. 260:2605-2608
- ORR and HOGAN, 1983, Pestic. Biochem. Biophys.
30 20:311-319
- PARKER and BLAKE, 1988, FEBS Letters 229:377-382
- PENG, MAYA and AYALA, 1986, PNAS USA 83:684-687
- POWLES (1984) Ann. Rev. Plant Physiol. 35: 15-44
- RABINOWITCH and FRIDOVICH, 1985, PLANTA 164: 524-528
- 35 - RABINOWITCH, SKLON and BUDOROSKI, 1982, Physiol. Plant.

54:369-374

- REISS, SPRENGEL and SCHALLER, 1984, EMBO J. 3:3317-3322
- ROBINSON, SMITH and GILLS, 1980, Plant Physiol. 65:755-759
- 5 - SACHS and HO, 1986, Ann. Rev. Plant. Physiol. 65:755-759
- SCHATZ, 1987, Eur. J. Biochem. 165:1-6
- SETO, HAYASHI and TENER, 1987, NAR 15:10601
- SHAALTIEL and GRESSEL, 1987, Plant Physiol. 85:869-871
- 10 - SIMOENS et al., 1986, NAR 14:8073-8090
- SINGLETON and SAINSBURY, 1987, Dictionary of Microbiology and Molecular Biology, Wiley and Sons
- SLATER, 1984, Meth.Enzymol. 105:283-293
- SNEDECOR and COCHRAN, 1980, "Statistical Methods", Iowa
- 15 State University Press.
- SOBEL and MARTINEZ, 1986, NAR 14:363-374
- STALLINGS, PATTRIDGE, STRONG and LUDWIG, 1985, J. Biol. Chem. 260:16424-16432
- STANSSENS, McKEOWN, FRIEDRICH and FRITZ, 1987, Manual
- 20 EMBO Laboratory course "Directed mutagenesis and protein engineering" held at Max Planck Institute für Biochemie, Martinsried, West-Germany, July 4-18, 1987.
- STEINITZ et al, 1979, Plant Science Letters p327-335
- STEINMAN, 1978, J. Biol. Chem. 253:8708-8720
- 25 - TAKAHASHI et al., 1985, PNAS USA 82:1931-1935
- TANAKA and SUGAHARA, 1980, Plant Cell Physiol. 21:601-612
- TANAKA, KONDO and SUGAHARA, 1982, Plant Cell Physiol. 23:999-1007
- 30 - VALVEKENS, VAN MONTAGU and VAN LYSEBETTENS, 1988, PNAS USA 85:5536-5540.
- VAN DEN BROECK, TIMKO, KAUSCH, CASHMORE, VAN MONTAGU and HERRERA-ESTRELLA, 1985, Nature, 358-363
- VELTEN and SCHELL, 1985, NAR 13:6998

35

48

- VELTEN, VELTEN, HAIN and SCHELL, 1984, EMBO J. 3:2723-2730
- VON HEIJNE, 1988, Bioch. Biophys. Acta 947: 307-333
- WATSON, 1984, NAR 12:5145-5164
- 5 - WINNACKER, 1987, From Genes to Clones, VCH Verlagsgesellschaft
- WU and GROSSMAN (eds.), 1987, "Methods in Enzymology", Vol. 153 Part D, Academic Press.
- 10 - YANNISH-PERRON, VIERA and MESSING, 1985, Gene 33:103-119
- ZACHEO, BLEVE-ZACHEO and LAMBERTI, 1982, Nemat. Medit. 10:75-80

15

20

25

30

35

C L A I M S

5 1. A recombinant SOD gene, preferably a chimaeric recombinant SOD gene, for transforming a plant cell, characterized by the following DNA fragments which are operably linked and in the same transcriptional unit:

10 a) an SOD gene encoding a metallo-superoxide dismutase, particularly a plant superoxide dismutase, quite particularly a Mn or Fe superoxide dismutase;

b) a promoter capable of directing expression, preferably overexpression, of said SOD gene in said plant cell; and

15 c) 3' transcription regulation signals for expression of said SOD gene in said plant cell.

20 2. The recombinant SOD gene of claim 1, further characterized by a targeting sequence which encodes a targeting peptide and which is located between said promoter and said SOD gene and is fused to, and in the same reading frame as, said SOD gene; said targeting peptide being adapted for translocation of said superoxide dismutase into mitochondria or chloroplasts of said cell or into the lumen of the endoplasmatic reticulum of said plant cell for secretion of said superoxide dismutase out of said plant cell.

25 3. The recombinant SOD gene of claim 1 or 2 wherein said SOD gene is a plant MnSOD gene or FeSOD gene, particularly a gene having the DNA sequence shown in Figure 1, 2, 3 or 5.

30 4. A plant cell, the genome, particularly the nuclear genome, of which is transformed with the recombinant SOD gene of anyone of claims 1-3.

35 5. A plant consisting of the plant cells of claim 4.

6. A plant cell culture consisting of the plant cells of claim 4.

7. A seed of the plant of claim 5.

8. Plant expression vectors characterized by the recombinant SOD gene of anyone of claims 1-3 between the border sequences of a disarmed T-DNA of Agrobacterium, particularly pEX1SOD, pEX3SOD, pEX4SOD, pEX5SOD, pEX6SOD, pEX7SOD or pEX8SOD.

9. A plant FeSOD' gene or MnSOD gene, particularly a gene having the DNA sequence shown in Figure 1, 2, 3 or 5.

10. A plant Fe superoxide dismutase or Mn superoxide dismutase, particularly a superoxide dismutase having the amino acid sequence shown in Figure 1, 2, 3 or 5.

11. In a process for producing plant and reproductive material, such as seeds, or for producing fruits of said plants including a foreign genetic material stably integrated in nuclear genome thereof and capable of being expressed therein as an RNA, protein or polypeptide, comprising the non-biological steps of: a) producing transformed plants cells or plant tissue including said foreign genetic material from starting plant cells or plant tissue not expressing said RNA, protein or polypeptide, b) producing regenerated plants or reproductive material of said plants or both from said transformed plant cells or plant tissue including said foreign genetic material, and c) optionally, biologically replicating said regenerated plants or reproductive material or both; wherein said step of producing said transformed plant cells or plant tissue including said foreign genetic material is characterized by: transforming the nuclear genome of said starting plant cells or plant tissue with a recombinant SOD gene of anyone of claims 1-3, as well as regulatory elements which are capable of enabling the expression of said foreign DNA sequence in




Fig. 14 B





Fig. 14 A


26/28

Fig. 13




25/26

Fig. 12 B



24/2B

Fig. 12A



22/28

Fig. 11 A






Fig. 10

21/28



20/26

Fig. 9 (cont.)



pEX5SOD

19/28


Fig. 9


Fig. 8 (cont.)

19/28

	Cys	Met	Asp	Leu	Gly	Leu	
(cp-tp)	TGC	ATG	GAT	CTG	GGC	TTG	(MnSOD)
	ACG	TAC	CTA	GAC	C	CGAAC	

pKAH1-SOD


BglII
BamHI
purify
fragment



MnSOD

pGSC1702

BamHI
cip



pEX4SOD

Fig. 8

17/02




Fig. 7 (cont.)





Fig. 7



14/28




Fig. 6

Figure 5 (continued)

13/28

660 670 680 690 700
AGTCAGTTCTAGGCTTAAAGCAGCAACAGCTTGAGCTGCTTAGCGAGAAG
V S S R L K A A T A

710 720 730 740 750
ACAGAAAGGAGGAAGAGGCCAAATCTAGCAGGCACGAGAGTAAATATTTGA

760 770 780 790 800
GACAGAATGATTTTTGTAAAGAGACACTATTTTCAATCCTGCTATCCTT

810 820 830 840 850
CTTTCTCAGTTGAGAATTTTAGATGTCTTATTATGTGCACTTTACTAGAG

860 870 880 890 900
AGTCAAGTGATGCTCTGTATTTGGAGGATAGTGTTATTTCTGTTCCTTAG

910 920 930 940 950
CAGCTGTTAATGGCAGGGAAAAATAATTCAAGTTGAGGTGTGGGACAACA

960 970 980 990 1000
ATGTAAGGACGTGAATAACAAATCTATTGCACCTTGTGCTTAATTIT

1010 1020 1030
AGAATTAGAATGAAAAAAACCCCCCCCCCCCCCCC

Figure 5

12/28

10 20 30 40 50
ATGACCATGATTACGAATTCGAGCTCGGTACCCGGGGATCCTCTAGAGTC
M T M I T N S S S V P G D P L E S

60 70 80 90 100
GACCTGCAGGGGGGGGGGGGGGGGGCTAAATTTGAACTCCAGCCTCCTCCTT
T C R G G G G A K F E L Q P P P

110 120 130 140 150
ATCCCATGGATGCTTTGGAGCCTCATATGAGTAGTAGAACGTTTGAATTC
Y P M D A L E P H M S S R T F E F

160 170 180 190 200
CACTGGGGGAAGCATCACAGGGCTTATGTGACAATTTAAACAAGCAAAT
H W G K H H R A Y V D N L N K Q I

210 220 230 240 250
AGACGGAAACAGAACTAGATGGAAAGACACTAGAAGACATAACTTGTTA
D G T E L D G K T L E D I I L V

260 270 280 290 300
CGTATAACAAAGGTGCTCCCTCCCGAGCATTCACAAATGCTGCTCAGGCC
T Y N K G A P L P A F N N A A Q A

310 320 330 340 350
TGGAATCATCAGTTTTTCTGGAATCAATGAAGCCCAACGGAGGAGGAGA
W N H Q F F W E S M K P N G G G E

360 370 380 390 400
GCCATCTGGTGAATTACTAGAACTAATCAACAGAGACTTTGGTTTCCTATG
P S G E L L E L I N R D F G S Y

410 420 430 440 450
ATGCATTTGTTAAAGAATTTAAGGCAGCTGCGGCAACACAATTTGGCTCT
D A F V K E F K A A A A T Q F G S

460 470 480 490 500
GGTTGGGCTGCTGCGATACAAACCTGAAGAGAAAAAGCTTGCCTTGGT
G W A W L A Y K P E E K K L A L V

510 520 530 540 550
GAAACTCCCAACGCTGAAAATCCTCTTGTGTTTGGGTTACACACCGCTCC
K T P N A E N P L V L G Y T P L

560 570 580 590 600
TCACCATAGACGTTTGGGAGCATGCTTACTATCTGGACTTTCAGAACCGG
L T I D V W E H A Y Y L D F Q N R

610 620 630 640 650
CGGCCTGACTACATATCTATCTTTATGGAGAGCTCGTGTCTGGGAAGC
R P D Y I S I F M E K L V S W E A

11/26

Figure 4 B (cont. 1)

```

710      720      730      740      750      760      770
AGTACTATTCCAGTATGGACCGATTCAAGGCTTGCTTCATAAACCAAGGCANGTAATAGAGATNGEAGTCT

780      790      800      810      820      830      840
CTAAAAAGGTAGTTCTACTGATCTTAAGGCCATGCCATGGAGTCTAAGATTCAAAATCGAGGATCTATCAG

850      860      870      880      890      900      910
AACTCGCCGTGAAGACTGGCGAACAGTTTCATACAGAGTCTTTTACGACTCAATCACAAGAGAAGAAATCTT

920      930      940      950      960      970      980
CGTCAACATGGTGGAGCACGACACTCTGGTCTACTCTCCMAAAATGTCAAAGATACAGTCTCAGAAGACCAA

990      1000     1010     1020     1030     1040     1050
AGGGCTATTGAGACTTTTCACAAAGGATAATTTCGGGAACCTCTCGGATTCATTTGCCAGCTAATCT

1060     1070     1080     1090     1100     1110     1120
GTCACTTCAGAAAGGACAGTAGAAAGGAAGGTGGCTCCTACAAATGCCCATCATTTGCGATPAAAGGAAA

1130     1140     1150     1160     1170     1180     1190
GGCTATCATTCAGATGCCCTCTGCGACAGTGGTCCCAAAGATGGACCCCAACCCAGGAGCATCGTG

1200     1210     1220     1230     1240     1250     1260
GAAAGAAGAGAGGTCCAAACCACCGTCTTCAAGCAAGTGGATTGATGTGACATCTCCACTGACCTAAGGG

1270     1280     1290     1300     1310     1320     1330
ATGACGCACAATCCCACTATCTCTTCGCAAGACCTTCTCTATATATAAGGAAGTTTCATTTTGGAGAG

1340     1350     1360     1370     1380
GACAGGCTGAATCACCAGTCTCTCTATATAAATCTATCTCTCTCTATATACCATGG

```

10/28

Figure 4 B

10 20 30 40 50 60 70
GAATCCAAATCCACCAAACTGAACCTAGCAGTTTCAGTTGCTCCTCTCAGAGACGAATCGGGTATTCA

80 90 100 110 120 130 140
ACACCTCATACCACTACTACGTGCTGTATTAACGGACCTCATGCGGGTATATATAGGATGACTGGGGTTGT

150 160 170 180 190 200 210
ACAAAGGCAGCAACAACGGTGTTCCTCCGGAGTTGGGCATAAGAAAGTTTGCCTACTATTACAGAGGCACAG

220 230 240 250 260 270 280
CAGCAGCTGACGGGTATACAACAAGTCAGCAACACAGATAGTTTGAACCTTCATCCCAAGGAGAGAGCTCA

290 300 310 320 330 340 350
ACTCAAGCCCAAGAGCTTTGCGAAGGCCCTAACAAAGCCCAACCAAGCAAAAGCCCACTGCTCAAGCTAG

360 370 380 390 400 410 420
GAACCAAAAGGCCAGCAGTGATCCAGCCCCCAAAAGGATCTCCCTTTGCCCGGAGATTACAAATGGACGA

430 440 450 460 470 480 490
TTTCCCTCTATCTTTACGATCTAGGAAGGAAGTTTCGAAGGTGAAGGTGACGACACTATGTTCCACCTGAT


500 510 520 530 540 550 560
AATGAGAAGGTTAGCCTCTTCGAATTTTCAGAAAGAATGCTGACCCACAGATGGTTAGAGAGGCTTAAGCAG

570 580 590 600 610 620 630
CAGGTCTCATCAAGACGATCTAACCAGTAACAATCTCCAGGAGATCAAAATACCTTCCCAAGAGGTTAA

640 650 660 670 680 690 700
AGATGCAGTCAAAAGATTACAGGACTAATTGTCATCAAGAACACAGAGAGAAAGACATATTTCTCAAGATCAGA

Fig. 4 A

9/28



8/28

Figure 3 (cont. 2)

675	702
GCA AAT TTG GTT CCT CTT CTG GGA ATA AAC GAT TGG GAA CAT GCA TAC TAC TTG	
Ala Asn Leu Val Pro Leu Leu Gly Ile Asp Val Trp Glu His Ala Tyr Tyr Leu	
729	756
CAG TAC AAA AAT GTA AGA CCT GAT TAT CTG AAG AAC ATA TGG AAA GTT ATG AAC	
Gln Tyr Lys Asn Val Arg Pro Asp Tyr Leu Lys Asn Ile Trp Lys Val Met Asn	
783	
TGG AAA TAT GCA AAT GAA GTT TAT GAG AAA GAA TGT CCT TGA	
Trp Lys Tyr Ala Asn Glu Val Tyr Glu Lys Glu Cys Pro	

7/28

Figure 3 (cont. 1)

CAT GAC GCC AAT TCC AAA GGA GAT GCT CCT ACC GTC GCC AAA TTG CAT AGC GCT	378
His Asp Ala Ile Ser Lys Gly Asp Ala Pro Thr Val Ala Lys Leu His Ser Ala	
351	
ATC AAA TTC AAC GGC GGA GGT CAC ATT AAC CAC TCG ATT TTC TGG AAG AAT CTT	432
Ile Lys Phe Asn Gly His Ile Asn His Ser Ile Phe Trp Lys Asn Leu	
405	
GCC CCT GTC CGC GAG GGT GGT GGT GAG CCT CCA AAG GGT TCT CTT GGN TGG GCT	486
Ala Pro Val Arg Glu Gly Gly Gly Glu Pro Pro Lys Gly Ser Leu Gly Trp Ala	
459	
ATC GAC ACT AAC TTT GGC TCC CTA GAA GCT TTA GTT CAA AAG ATG AAT GCA GAA	540
Ile Asp Thr Asn Phe Gly Ser Leu Glu Ala Leu Val Gln Lys Met Asn Ala Glu	
513	
GGT GCT GCT TTA CAG GGC TCT GGC TCT GGC TGG GTG TGG CTT GGT GTG GAC AAA GAG CTT	594
Gly Ala Ala Leu Leu Gln Gly Ser Gly Trp Val Trp Leu Gly Val Asp Lys Glu Leu	
567	
AAG CGC CTG GTG ATT GAA ACC ACT GCT AAT CAG GAC CCT TTG GTT TCT AAA GGA	648
Lys Arg Leu Val Ile Glu Thr Thr Ala Asn Gln Asp Pro Leu Val Ser Lys Gly	
621	

6/28

Figure 3

ATG GCT TCT ATG ATA TCC TCT TCC GCT GTG ACA ACA GTC AGC CGT GCC TCT AGG	27	54
MET Ala Ser MET Ile Ser Ser Ser Ala Val Thr Thr Val Ser Arg Ala Ser Arg		
GGG CAA TCC GGC GCA GTG GCT CCA TTC GGC GGC CTC AAA TCC ATG ACT ACT GGA TTC	81	108
Gly Gln Ser Ala Ala Val Ala Pro Phe Gly Gly Leu Lys Ser MET Thr Gly Phe		
CCA GTG AAG AAG GTC AAC ACT GAC ATT ACT TCC ATT ACA AGC AAT GGT GGA AGA	135	162
Pro Val Lys Lys Val Asn Thr Asp Ile Thr Ser Ile Thr Ser Asn Gly Gly Arg		
GTA AAG TGC ATG GAT CTG GGC TTG CAG ACC TTT TCG CTC CCC GAT CTC CCC TAC	↓ 189	216
Val Lys Cys MET ASP Leu Gly Leu Gln Thr Phe Ser Leu Pro Asp Leu Pro Tyr		
GAC TAT GGA GCA CTG GAG CCG GCA ATT AGC GGT GAC ATA ATG CAG CTC CAC CAC	243	270
ASP Tyr Gly Ala Leu Gln Pro Ala Ile Ser Gly Asp Ile MET Gln Leu His His		
CAG AAT CAC CAT CAG ACT TAC GTC ACC AAT TAC AAT AAA GCC CTT GAA CAG CTA	297	324
Gln Asn His His Gln Thr Tyr Val Thr Asn Tyr Asn Lys Ala Leu Gln Gln Leu		

Figure 2 (cont.)

351

AAC	TTT	GGC	TCC	CYA	GAA	GCT	TTA	GTT	CAA	AAG	ATG	AAT	GCA	GAA	GGT	GCT	GCT
ASN	Phe	Gly	Ser	Leu	Glu	Ala	Leu	Val	Gln	Lys	Met	Asn	Ala	Gln	Gly	Ala	Ala

378

405
TTTA CAG GGC TCT GGC TGG GTG TGG CTT GAC AAC GAG CTT AAG CGC CTS
Leu Glu Gly Ser Gly Tyr Val Tyr Leu Gly Val Asp Lys Glu Leu Lys Arg Leu
432

459
GTG ATT GAA ACC ACT GCT AAT CAG GAC CCT TTG GTP TCT AAA GGA GCA AAT TTG
Val Ile Glu Thr Thr Ala Asn Glu Asp Pro Leu Val Ser Lys Gly Ala Asn Leu

GTG	CCT	CTG	GGG	ATA	GAC	GTT	TGG	GAA	CAT	GCA	TAC	TAC	TGG	CAG	TAC	AAA	540
val	pro	leu	leu	gly	ile	asp	val	trp	glu	his	ala	tyr	tyr	leu	glu	tyr	lys

AAU	GTA	AGA	CCT	GAT	TAT	CTG	AGG	AAC	ATA	TGG	AAA	GTT	ATG	AAC	TGG	AAA	TAT	594
AGA	VAL	ATG	PRO	ASP	TYR	LEU	LVS	ASN	ILE	TRP	LYG	VAL	NET	ASN	TRP	LVS	TYT	

621
GCA AAT GAA GTT TAT GAG AAA GAA TGT CCT TGA
Ala Asn Glu Val Tyr Glu Lys Glu Cys Pro .

4/28

Figure 2

ATG GAT CTG GGC TTG CAG ACC TTT TCG CTC CCC GAT CTC CCC TAC GAC TAT GGA	27	54
MET ASP LEU GLY LEU GLN THR PHE SER LEU PRO ASP LEU PRO TYR ASP TYR GLY		
GCA CTG CAG CCG GCA ATT AGC GGT GAC ATA ATG CAG CTC CAC CAG CAG AAT CAC	81	108
ALA LEU GLU PRO ALA ILE SER GLY ASP ILE MET GLN LEU HIS HIS GLN ASN HIS		
CAT CAG ACT TAC GTC ACC AAT TAC AAT AAA GCC CTT GAA CAG CTA CAT GAC GCC	135	162
HIS GLN THR TYR VAL THR ASN TYR ASN LYS ALA LEU GLU GLN LEU HIS ASP ALA		
ATT TCC AAA GGA GAT GCT CCT CCT ACC GTC GCC AAA TTG CAT AGC GCT ATC AAA TTC	189	216
ILE SER LYS GLY ASP ALA PRO THR VAL ALA LYS LEU HIS SER ALA ILE LYS PHE		
AAC GGC GGA GGT CAC ATT AAC CAC TCG ATT TTC TGG AAG AAT CTT GCC CCT GTC	243	270
ASN GLY GLY GLY HIS ILE ASN HIS SER ILE PHE TRP LYS ASN LEU ALA PRO VAL		
CGC GAG GGT GGT GGT GAG CCT CCA AAG GGT TCT CTT GGT TGG GCT ATC GAC ACT	297	324
ARG GLU GLY GLY GLY GLU PRO PRO LYS GLY SER LEU GLY TRP ALA ILE ASP THR		

Figure 1 (cont. 2)

TAC AAA AAT GTA AGA CCT GAT TAT CTG AAG AAC ATA TGG AAA GTT ATG AAC TGG	648
Tyr Lys Asn Val Arg Phe Asp Tyr Leu Lys Asn Ile Tyr Lys Val Met Asn Tyr	
AAA TAT GCA AAT GAA GTT TAT GAG AAA GAA TGT CCT TGAACAGGGA TATTGATGT	
Lys Tyr Ala Asn Glu Val Tyr Glu Lys Glu Cys Phe	
TGTTTTCAGG ACGTCGTAA AACTTTTGA TGGGAAATAA GGCTGAGTGA CATGAGCAGG	
TGTCCTGTTT TTCTTTCGATG TAGTCGCTGG CTGATGTACT TGATGTATTT CTGGAAAAGG	
TTGATGTATG TACTTGTAT ATGGAGGCTA AATAAACA CTCTATCGTT TGAGCGGAAA	
CCCCCCCCCC CCCC	

*

675

2/28

Figure 1 (cont. 1)

AAA TTC AAC GGC GGA GGT CAC ATT AAC CAC TCG ATT TTC TGG AAG AAT CTT GGC	324
Lys Phe Asn Gly Gly His Ile Asn His Ser Ile Phe Tip Lys Asn Leu Ala	297
CCT GTC CGC GAG GGT GGT GGT CCA AAG GGT TCT CTT GGT TGG GCT ATC	378
Pro Val Arg Glu Gly Gly Gly Pro Pro Lys Gly Ser Leu Gly Tip Ala Ile	351
GAC ACT AAC TTT GGC TCC CTA GAA GCT TTA GTT CAA AAG ATG AAT GCA GAA GGT	432
Asp Thr Asn Phe Gly Ser Leu Glu Ala Leu Val Gln Lys Met Asn Ala Glu Gly	405
GCT GCT TTA CAG GGC TCT GGC TGG TGG TGG CTT GGT GTG GAC AAA GAG CTT AAG	486
Ala Ala Leu Gln Gly Ser Gly Trp Val Tip Lys Gly Val Asp Lys Glu Leu Lys	459
CGC CTG GTG ATT GAA ACC ACT GCT ACT GCT AAT CAG GAC CCT TTG GTT TCT AAA GGA GCA	540
Arg Leu Val Ile Glu Thr Thr Ala Asn Gln Asp Pro Leu Val Ser Lys Gly Ala	513
AAT TTG GTT CCT CTT CTG GGA ATA GAC GTT TGG GAA CAT GCA TAC TAC TTG CAG	594
Asn Leu Val Pro Leu Leu Gly Ile Asp Val Trp Glu His Ala Tyr Tyr Leu Gln	567

1/28

Figure 1

GGGGGGGGGG GGGGGGCTGG CCGCTCTGGG CATGACCTGC AACTATATAAA GGACACCCATA

GAGTTAACAG CTAGAAAGCA TTTAGGAATA TCTCAAAA

27
 54
 ATG GCA CTA CGA ACC CTA GNG ACC AGA CCG ACC TTA GCA ACA GGG CTA GGG TTC
 MET Ala Leu Arg Thr Leu Val Ser Arg Arg Thr Leu Ala Thr Gly Leu Gly Phe

81
 108
 CGC CAG CAA CTC CGC GGC TTG CAG ACC TTT TCG CTC CCC GAT CTC CCC TAC GAC
 Arg Gln Gln Leu Arg Gly Leu Gln Thr Phe Ser Leu Pro Asp Leu Pro Tyr Asp

135
 162
 TAT GGA GCA CTG GAG CCG GCA ATT AGC GGT GAC ATA ATG CAG CTC CAC CAC CAG
 Tyr Gly Ala Leu Gln Pro Pro Ala Ile Ser Gly Asp Ile MET Gln Leu His His Gln

189
 216
 AAT CAC CAT CAG ACT TAC GTC ACC AAT TAC AAT AAA GCC CTT GAA CAG CTA CAT
 Asn His His Gln Thr Tyr Val Thr Asn Tyr Asn Lys Ala Leu Gln Leu His

243
 270
 GAC GCC ATT TCC AAA GGA GAT GGT CCT ACC GTC GCC AAA TTG CAT AGC GGT ATC
 Asp Ala Ile Ser Lys Gly Asp Ala Pro Thr Val Ala Lys Leu His Ser Ala Ile

EP 8901025
SA 30815

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A- 0213628	11-03-87	AU-A- 6219186 JP-A- 62115277	05-03-87 26-05-87

64501 WORM 10479

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
	and plastocyanin transit peptides", pages 377-388, see the whole article -----	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
P,X	The EMBO Journal, vol. 8, no. 1, 1989, IRL Press, C. Bowler et al.: "The induction of manganese superoxide dismutase in response to stress in Nicotiana plumbaginifolia", pages 31-38, see the whole article --	9,10
A	EP, A, 0213628 (YEDA RESEARCH) 11 March 1987, see page 14, lines 3-7; claim 13 --	1-12
A	EMBL Sequence Database Entry, accession number X12540,- "Maize mRNA for superoxide dismutase-3 isoenzyme (EC 1.15.1.1) --	9,10
A	Chemical Abstracts, vol. 107, 1987, (Columbus, Ohio, US), J.M. Palma et al.: "Increased levels of peroxisomal active oxygen-related enzymes in copper-tolerant pea plants", page 259, abstract no. 230884y, Plant Physiol. 1987, 85(2), 570-4, see the abstract --	3,12
A	Chemical Abstracts, vol. 106, 1987 (Columbus, Ohio, US), G.L. Matters: "Influence of environ- mentally induced oxidative stress on the expression of catalase and super- oxide dismutase genes in Zea mays L", page 168, abstract no. 44954a, Diss. Abstr. Int. B 1986, 47(6), 2295, see the abstract --	12
A	Proc. Natl. Acad. Sci. USA, vol. 84, January 1987, R.E. Cannon et al.: "Cloning of cDNA for maize superoxide dismutase 2 (SOD2)", pages 179-183, see the abstract; introduction --	1-12
A	Plant Molecular Biology, vol. 9, 1987, Martinus Nijhoff Publishers (Dordrecht, NL), S. Smeekeens et al.: "Import into chloroplasts of a yeast mitochondrial protein directed by ferredoxin and ./.	1-12

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
Y	NH ₂ -terminal amino acid-sequence analysis of proteins electroblotted from two-dimensional gel-separated total extracts", pages 4806-4810, see page 4810, paragraph "Note added in proof"	3,9
O,X	-- Biological Abstracts/RRM, BR 33:64036, J. White et al.: "The compartmentalization of the maize manganese superoxide dismutase SOD-3", & Annual Meeting of the American Society of Plant Physiologists, St. Louis, Missouri, USA, 19-23 July 1987. Plant Physiol, (Bethesda) 83 (4 suppl.), 1987, 106, see the abstract	9,10
X	-- Chemical Abstracts, vol. 107, 1987, (Columbus, Ohio, US), J.A. White et al.: "In vitro synthesis and processing of manganese-superoxide dismutase (SOD-3) into maize mitochondria", page 391. abstract no. 232177u, & Biochim. Biophys. Acta 1987, 926(1), 16-25, see the abstract	9,10
X	-- Eur. J. Biochem., vol. 146, 1985, FEBS, J. Kwiatowski et al.: "Isolation and characterization of an iron-containing superoxide dismutase from tomato leaves, Lycopersicon esculentum", pages 459-466, see the whole article	10
Y	--	9
Y	Chemical Abstracts, vol. 107, 1987, (Columbus, Ohio, US), Y. Shaaltiel et al.: "Biochemical analysis of paraquat resistance in Conyza leads to pinpointing synergists for oxidant generating herbicides", page 273, abstract no. 192885q, & Pestic. Sci. Biotechnol., Proc. Int. Congr. Pestic. Chem., 6th 1986 (Pub. 1987) 183-6, see abstract	2

INTERNATIONAL SEARCH REPORT

International Application No PCT/EP 89/01025

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) * According to International Patent Classification (IPC) or to both National Classification and IPC IPC ⁵ : C 12 N 15/53, C 12 N 15/82, C 12 N 5/10, A 01 H 5/00, A 01 H 5/10, C 12 N 9/02		
II. FIELDS SEARCHED <div style="display: flex; justify-content: space-between;"> Classification System Minimum Documentation Searched⁷ </div> <div style="display: flex; justify-content: space-between;"> IPC⁵ C 12 N, A 01 H </div> <div style="display: flex; justify-content: space-between;"> Classification Symbols </div>		
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched *		
III. DOCUMENTS CONSIDERED TO BE RELEVANT*		
Category ⁸	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	Genetic Technology News, vol. 6, no. 8, August 1986, A.S. Hester: "Recombinant bacteria and plants for agriculture at AGS", pages 5,8, see page 5, last 2 lines - page 8, lines 1-3	1,4,5,6,11, 12
Y	--	2,3,7,8
X	Derwent WPIL, accession no. 84-312582, Y. Grower: "DNA coding for human or plant super:oxide dismutase - useful e.g. for diagnosis of Downs syndrome", & AU, A, 8427461 (YEDA RESEARCH & DEVELOPMENT CO. LTD) 1 November 1984, see the abstract	1,4,5,6,11, 12
Y	--	2,3,7,8
X	Proc. Natl. Acad. Sci. USA, vol. 84, July 1987, G. Bauw et al.: "Alterations in the phenotype of plant cells studied by ./.	9,10
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>* Special categories of cited documents:¹⁴</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search <div style="text-align: center;">30th November 1989</div>		Date of Mailing of this International Search Report <div style="text-align: center;">03 DEC 1989</div>
International Searching Authority <div style="text-align: center;">EUROPEAN PATENT OFFICE</div>		Signature of Authorized Officer <div style="text-align: center;">C.D. v.d. Vliet </div>

Form PCT/ISA/210 (second sheet) (January 1985)




Fig. 14 B





Fig. 14A


26/28

Fig. 13




25/28

Fig. 12 B



24/28

Fig. 12A



23/28

Fig. 11 B




Fig. 11 A






Fig. 10

21/28



20/28

Fig. 9 (cont.)**pEX5SOD**

19/28


Fig. 9



Fig. 8 (cont.)

18/28

	Cys	Met	Asp	Leu	Gly	Leu	
(cp-tp)	TGC	ATG	GAT	CTG	GGC	TG	(MnSOD)
	ACG	TAC	CTA	GAC	C	CGAAC	

pKAH1-SOD

BglII
BamHI
purify
fragment



pEX4SOD


MnSOD

pGSC1702

BamHI
cip

Fig. 8

17/28



16/28

Fig. 7 (cont.)





Fig. 7



14/28




Fig. 6

Figure 5 (continued)

13/28

```
      660      670      680      690      700
AGTCAGTTCTAGGCTTAAAGCAGCAACAGCTTGAGCTGCTTAGCGAGAAG
  V  S  S  R  L  K  A  A  T  A

      710      720      730      740      750
ACAGAAAGGAGGAAGAGGCCAAATCTAGCAGGCACGAGAGTAAATATTTGA

      760      770      780      790      800
GACAGAATGATTTTTGTAAAGAGACACTATTTTCAATCCTGCTATCCTT

      810      820      830      840      850
CTTTCTCAGTTGAGAATTTTAGATGTCTTATTATGTGCACTTTACTAGAG

      860      870      880      890      900
AGTCAAGTGATGCTCTGTATTTGGAGGATAGTGTTATTTCTGTTCTTTAG

      910      920      930      940      950
CAGCTGTTAATGGCAGGGAAAAATAATTCAAGTTGAGGTGTGGGACAACA

      960      970      980      990     1000
ATGTAAGGACGTGAATAAACAAATCTATTGCACTTTGGTGCCCTAATTTT

     1010     1020     1030
AGAATTAGAATGAAAAAAAAACCCCCCCCCCCCCCCC
```

Figure 5

12/28

10 20 30 40 50
ATGACCATGATTACGAATTCGAGCTCGGTACCCGGGGATCCTCTAGAGTC
M T M I T N S S S V P G D P L E S

60 70 80 90 100
GACCTGCAGGGGGGGGGGGGGGGCTAAATTTGAACTCCAGCCTCCTCCTT
T C R G G G G A K F E L Q P P P

110 120 130 140 150
ATCCCATGGATGCTTTGGAGCCTCATATGAGTAGTAGAACGTTTGAATTC
Y P M D A L E P H M S S R T F E F

160 170 180 190 200
CACTGGGGGAAGCATCACAGGGCTTATGTGACAATTTAAACAAGCAAAT
H W G K H H R A Y V D N L N K Q I

210 220 230 240 250
AGACGGAACAGAACTAGATGGAAAGACACTAGAAGACATAATACTTGTTA
D G T E L D G K T L E D I I L V

260 270 280 290 300
CGTATAACAAAGGTGCTCCCCTCCAGCATTCAACAATGCTGCTCAGGCC
T Y N K G A P L P A F N N A A Q A

310 320 330 340 350
TGGAATCATCAGTTTTTCTGGGAATCAATGAAGCCCAACGGAGGAGGAGA
W N H Q F F W E S M K P N G G G E

360 370 380 390 400
GCCATCTGGTGAATTACTAGAACTAATCAACAGAGACTTTGGTTCCTATG
P S G E L L E L I N R D F G S Y

410 420 430 440 450
ATGCATTTGTTAAAGAATTTAAGGCAGCTGCGGCAACACAATTTGGCTCT
D A F V K E F K A A A A T Q F G S

460 470 480 490 500
GGTTGGGCCTGGCTCGCATACAAACCTGAAGAGAAAAAGCTTGCCTTGGT
G W A W L A Y K P E E K K L A L V

510 520 530 540 550
GAAAACTCCCAACGCTGAAAATCCTCTTGTGTTTGGGTTACACACCGCTCC
K T P N A E N P L V L G Y T P L

560 570 580 590 600
TCACCATAGACGTTTGGGAGCATGCTTACTATCTGGACTTTCAGAACCGG
L T I D V W E H A Y Y L D F Q N R

610 620 630 640 650
CGGCCTGACTACATATCTATCTTTATGGAGAAGCTCGTGTTCGTGGGAAGC
R P D Y I S I F M E K L V S W E A

11/28

Figure 4 B (cont. 1)

710 720 730 740 750 760 770
AGTACTATTCCAGTATGGACGATTCAAGGCTTGCTTCATAAACCAAGCAAGTAATAGAGATTGGAGTCT

780 790 800 810 820 830 840
CTAAAGGTAGTTCCCTACTGAATCTAAGGCCATGCATGGAGTCAAGATTCAAAATCGAGGATCTAACAG

850 860 870 880 890 900 910
AACTCGCCGTGAAGPCTGGCGAACACAGTTTCATACAGAGTCTTTTACGACTCAATGACAAGAAAAATCTT

920 930 940 950 960 970 980
CGTCAACATGGTGGAGCACGACACTCTGGTCTACTCCAAAAATGTCAAAGATACAGTCTCAGAAAGACCAA

990 1000 1010 1020 1030 1040 1050
AGGGCTATTGAGACTTTTCAACAAAGGATAATTTCGGGAAACCTCTCGGATTCCATTGCCCAGCTATCT

1060 1070 1080 1090 1100 1110 1120
GTCACCTTCGAAAGGACACGTAGAAAAGGAAGGTGGTCCCTACAAAATGCCATCATTCGGATAAAGGAAA

1130 1140 1150 1160 1170 1180 1190
GGCTATCATTAAGATGCCCTCTGCCGACAGTGGTCCCAAGATGGACCCCAAGAGGAGCATCGTG

1200 1210 1220 1230 1240 1250 1260
GAAAAAGAGACGTTCCAAACCACGCTTTCAAAGCAAGTGGATTGATGTGACATCTCCACTGACGTAAGGG

1270 1280 1290 1300 1310 1320 1330
ATGACGCACAATCCCACATATCCTTCGCAAGACCCCTTCCTCTATATATAAGGAAGTTCAATTTTCATTTGGAGAG

1340 1350 1360 1370 1380
GACACGCTGAAATCACCAGTCTCTCTATATAAATCTATCTCTCTCTCTATAACCATGG

10/28

Figure 4 B

10 20 30 40 50 60 70
GAATTCCAATCCCAACCAAAACCTGAACCTAGCAGTTCA GTTGCTCCTCTCAGAGACGAATCGGGTATTCA

80 90 100 110 120 130 140
ACACCCCTCATACCAACTACTACGTCGTGTATTAACGGACCTCATGCCGGTATATACGATGACTGGGGTTGT

150 160 170 180 190 200 210
ACAAAGGCAGCAACAACGGTGTTCCTCCGGAGTTGGGCATAGAAGTTTGCCACTATTACAGAGGCAAGAG

220 230 240 250 260 270 280
CAGCAGCTGACGCGGTATACAACAAGTCAGCAACACAGATAGGTGAACCTTCATCCCCAAAGGAGAAGCTCA

290 300 310 320 330 340 350
ACTCAAGCCCCAAGAGCTTTGCCGAAGGCCCTTAACAAGCCCCACCAAGCAAAAAGCCCCACTGCTCACGCTAG

360 370 380 390 400 410 420
GAACCAAAAGGGCCAGCAGTGATCCAGCCCCCAAAAGAGATCTCCTTTGCCCCGGAGATTACAATGGACCA

430 440 450 460 470 480 490
TTTTCCTCTATCTTTACGATCTTAGGAAGGAAGTTCGAAGGTGAAGGTGACGACACTATGTTTCAACCACCTGAT


500 510 520 530 540 550 560
AATGAGAAGGTTAGCCCTCTTCAATTTCAAGAAAGAAATGCTGACCCACAGATGGTTAGAGAGGCCCTACGCAG

570 580 590 600 610 620 630
CAGGTCTCATCAAGACGATCTACCCGAGTAACAATCTCCAGGAGATCAAATACCTTCCCAAGAAGGTTAA

640 650 660 670 680 690 700
AGATGCAGTCAAAAAGATTGAGGACTAATTGCATCAAGAACAACAGAGAGAAAGACATATTTCTCAAGATCAGA

Fig. 4 A

9/28



8/28

Figure 3 (cont. 2)

GCA AAT TTG GTT CCT CTT CTG GGA ATA GAC GTT TGG GAA CAT GCA TAC TAC TTG	702
Ala Asn Leu Val Pro Leu Leu Gly Ile Asp Val Trp Glu His Ala Tyr Tyr Leu	
CAG TAC AAA AAT GTA AGA CCT GAT TAT CTG AAG AAC ATA TGG AAA GTT ATG AAC	756
Gln Tyr Lys Asn Val Arg Pro Asp Tyr Leu Lys Asn Ile Trp Lys Val MET Asn	
TGG AAA TAT GCA AAT GAA GTT TAT GAG AAA GAA TGT CCT TGA	783
Trp Lys Tyr Tyr Ala Asn Glu Val Tyr Glu Lys Glu Cys Pro	

7/28

Figure 3 (cont. 1)

CAT GAC GCC ATT TCC AAA GGA GAT GCT	351	CCT ACC GTC GCC AAA TTG CAT AGC GCT	378
His Asp Ala Ile Ser Lys Gly Asp Ala Pro		Thr Val Ala Lys Leu His Ser Ala	
ATC AAA TTC AAC GGC GGA GGT CAC ATT AAC	405	CAC TCG ATT TTC TGG AAG AAT CTT	432
Ile Lys Phe Asn Gly His Gly Gly Ile Asn		His Ser Ile Phe Trp Lys Asn Leu	
GCC CCT GTC CGC GAG GGT GGT GGT GAG	459	CCT CCA AAG GGT TCT CTT GGT TGG GCT	486
Ala Pro Val Arg Glu Gly Gly Gly Glu Pro		Pro Lys Gly Ser Leu Gly Trp Ala	
ATC GAC ACT AAC TTT GGC TCC CTA GAA	513	GCT TTA GTT CAA AAG ATG AAT GCA GAA	540
Ile Asp Thr Asn Phe Gly Ser Leu Glu Glu		Ala Leu Val Gln Lys MET Asn Ala Glu	
GGT GCT GCT TTA CAG GGC TCT GGC TGG	567	TGG GTG TGG CTT GGT GAC AAA GAG CTT	594
Gly Ala Ala Leu Gln Gly Ser Gly Trp Trp		Val Trp Leu Gly Val Asp Lys Glu Leu	
AAG CGC CTG GTG ATT GAA ACC ACT GCT	621	AAT CAG GAC CCT TTG GTT TCT AAA GGA	648
Lys Arg Leu Val Ile Glu Thr Thr Ala Asn		Gln Asp Pro Leu Val Ser Lys Gly	

6/28

Figure 3

ATG GCT TCT ATG ATA TCC TCT TCC GCT GTG ACA ACA GTC AGC CGT GCC TCT AGG	54
MET Ala Ser MET Ile Ser Ser Ser Ala Val Thr Thr Val Ser Arg Ala Ser Arg	
GGG CAA TCC GCC GCA GTG GCT CCA TTC GGC GGC CTC AAA TCC ATG ACT GGA TTC	108
Gly Gln Ser Ala Ala Val Ala Pro Phe Gly Gly Leu Lys Ser MET Thr Gly Phe	
CCA GTG AAG AAG GTC AAC ACT GAC ATT ACT TCC ATT ACA AGC AAT GGT GGA AGA	162
Pro Val Lys Lys Val Asn Thr Asp Ile Thr Ser Ile Thr Ser Asn Gly Gly Arg	
GTA AAG TGC ATG GAT CTG GGC TTG CAG ACC TTT TCG CTC CCC GAT CTC CCC TAC	216
Val Lys Cys MET Asp Leu Gly Leu Gln Thr Phe Ser Leu Pro Asp Leu Pro Tyr	
GAC TAT GGA GCA CTG GAG CCG GCA ATT AGC GGT GAC ATA ATG CAG CTC CAC CAC	270
Asp Tyr Gly Ala Leu Glu Pro Ala Ile Ser Gly Asp Ile MET Gln Leu His His	
CAG AAT CAC CAT CAG ACT TAC GTC ACC AAT TAC AAT AAA GCC CTT GAA CAG CTA	324
Gln Asn His His Gln Thr Tyr Val Thr Asn Tyr Asn Lys Ala Leu Glu Gln Leu	

5/28

Figure 2 (cont.)

AAC TTT GGC TCC CTA GAA GCT TTA GTT CAA AAG ATG AAT GCA GAA GGT GCT GCT	378
Asn Phe Gly Ser Leu Glu Ala Leu Val Gln Lys MET Asn Ala Glu Gly Ala Ala	
TTA CAG GGC TCT GGC TGG GTG TGG CTT GGT GTG GAC AAA GAG CTT AAG CGC CTG	432
Leu Gln Gly Ser Gly Trp Val Trp Leu Gly Val Asp Lys Glu Leu Lys Arg Leu	
GTG ATT GAA ACC ACT GCT AAT CAG GAC CCT TTG GTT TCT AAA GGA GCA AAT TTG	486
Val Ile Glu Thr Thr Ala Asn Gln Asp Pro Leu Val Ser Lys Gly Ala Asn Leu	
GTT CCT CTT CTG GGA ATA GAC GTT TGG GAA CAT GCA TAC TAC TTG CAG TAC AAA	540
Val Pro Leu Leu Gly Ile Asp Val Trp Glu His Ala Tyr Tyr Leu Gln Tyr Lys	
AAT GTA AGA CCT GAT TAT CTG AAG AAC ATA TGG AAA GTT ATG AAC TGG AAA TAT	594
Asn Val Arg Pro Asp Tyr Leu Lys Asn Ile Trp Lys Val MET Asn Trp Lys Tyr	
GCA AAT GAA GTT TAT GAG AAA GAA TGT CCT TGA	621
Ala Asn Glu Val Tyr Glu Lys Glu Cys Pro	

4/28

Figure 2

54
 ATG GAT CTG GGC TTG CAG ACC TTT TCG CTC CCC GAT CTC CCC TAC GAC TAT GGA
 MET Asp Leu Gly Leu Gln Thr Phe Ser Leu Pro Asp Leu Pro Tyr Asp Tyr Gly

81
 GCA CTG GAG CCG GCA ATT AGC GGT GAC ATA ATG CAG CTC CAC CAC GAG AAT CAC
 Ala Leu Glu Pro Ala Ile Ser Gly Asp Ile MET Gln Leu His His Gln Asn His

135
 CAT CAG ACT TAC GTC ACC AAT TAC AAT AAA GCC CTT GAA CAG CTA CAT GAC GCC
 His Gln Thr Tyr Val Thr Asn Tyr Asn Lys Ala Leu Glu Gln Leu His Asp Ala

189
 ATT TCC AAA GGA GAT GCT CCT ACC GTC GCC AAA TTG CAT AGC GCT ATC AAA TTC
 Ile Ser Lys Gly Asp Ala Pro Thr Val Ala Lys Leu His Ser Ala Ile Lys Phe

243
 AAC GGC GGA GGT CAC ATT AAC CAC TCG ATT TTC TGG AAG AAT CTT GCC CCT GTC
 Asn Gly Gly Gly His Ile Asn His Ser Ile Phe Trp Lys Asn Leu Ala Pro Val

297
 CGC GAG GGT GGT GGT GAG CCT CCA AAG GGT TCT CTT GGT TGG GCT ATC GAC ACT
 Arg Gln Gly Gly Gly Glu Pro Pro Lys Gly Ser Leu Gly Trp Ala Ile Asp Thr

3/28

Figure 1 (cont. 2)

	621		648
TAC AAA AAT GTA AGA CCT GAT TAT CTG AAG AAC ATA TGG AAA GTT ATG AAC TGG			
Tyr Lys Asn Val Arg Pro Asp Tyr Leu Lys Asn Ile Trp Lys Val MET Asn Trp			
		*	
AAA TAT GCA AAT GAA GTT TAT GAG AAA GAA TGT CCT TGAACAGGGA TATTTGATGT	675		
Lys Tyr Ala Asn Glu Val Tyr Glu Lys Glu Cys Pro			
TGTTTGGAGG ACGTCTGTAA AACTTTTGA TGGGAAATAA GGCTGAGTGA CATGAGCAGG			
TGTCCTGTTF TTCTTGCATG TAGTCGCTGG CTGATGTATT TGATGTATTT CTGGAAAAGG			
TTGATGTATG TACTTGATAT ATGGAGCCCTA AATAAACTA CTCATATCGTT TGAGCGCAAA			
CCCCCCCCCC CCCC			

2/28

Figure 1 (cont. 1)

AAA TTC AAC GGC GGA GGT CAC ATT AAC CAC TCG ATT TTC TGG AAG AAT CTT GCC	324
Lys Phe Asn Gly Gly His Ile Asn His Ser Ile Phe Trp Lys Asn Leu Ala	297
CCT GTC CGC GAG GGT GGT GGT CCA AAG GGT TCT CTT GGT TGG GCT ATC	378
Pro Val Arg Glu Gly Gly Glu Pro Pro Lys Gly Ser Leu Gly Trp Ala Ile	351
GAC ACT AAC TTT GGC TCC CTA GAA GCT TTA GTT CAA AAG ATG AAT GCA GAA GGT	432
Asp Thr Asn Phe Gly Ser Leu Glu Ala Leu Val Gln Lys MET Asn Ala Glu Gly	405
GCT GCT TTA CAG GGC TCT GGC TGG GTG TGG CTT GGT GTG GAC AAA GAG CTT AAG	486
Ala Ala Leu Gln Gly Ser Gly Trp Val Trp Leu Gly Val Asp Lys Glu Leu Lys	459
CGC CTG GTG ATT GAA ACC ACT GCT ACT GCT AAT CAG GAC CCT TTG GTT TCT AAA GGA GCA	540
Arg Leu Val Ile Glu Thr Thr Ala Asn Gln Asp Pro Leu Val Ser Lys Gly Ala	513
AAT TTG GTT CCT CTT CTG GGA ATA GAC GTT TGG GAA CAT GCA TAC TAC TTG CAG	594
Asn Leu Val Pro Leu Leu Gly Ile Asp Val Trp Glu His Ala Tyr Tyr Leu Gln	567

1/28

Figure 1

GGGGGGGGG GGGGGGCTGG CCTCTCTGGG CATGACCTGC AACTATAAAA GGACACCATA

GAGTTAACAG CTAGAAAGCA TTTAGGAATA TCTCAAAA

27
 ATG GCA CTA CGA ACC CTA GTG AGC AGA CGG ACC TTA GCA ACA GGG CTA GGG TTC
 MET Ala Leu Arg Thr Leu Val Ser Arg Arg Thr Leu Ala Thr Gly Leu Gly Phe

↓
 81
 CGC CAG CAA CTC CGC GGC TTG CAG ACC TTT TCG CTC CCC GAT CTC CCC TAC GAC
 Arg Gln Gln Leu Arg Gly Leu Gln Thr Phe Ser Leu Pro Asp Leu Pro Tyr Asp

135
 TAT GGA GCA CTG GAG CCG GCA ATT AGC GGT GAC ATA ATG CAG CTC CAC CAC CAG
 Tyr Gly Ala Leu Glu Pro Ala Ile Ser Gly Asp Ile MET Gln Leu His His Gln

189
 AAT CAC CAT CAG ACT TAC GTC ACC AAT TAC AAT AAA GCC CTT GAA CAG CTA CAT
 Asn His His Gln Thr Tyr Val Thr Asn Tyr Asn Lys Ala Leu Glu Gln Leu His

243
 GAC GCC ATT TCC AAA GGA GAT GCT CCT ACC GTC GCC AAA TTG CAT AGC GCT ATC
 Asp Ala Ile Ser Lys Gly Asp Ala Pro Thr Val Ala Lys Leu His Ser Ala Ile

51

said plant cells or plant tissue, to cause the stable integration of said foreign DNA sequence in transformed plant cells or plant tissue, as well as in plants and reproductive material produced therefrom throughout subsequent generations.

12. A method for rendering a plant more resistant or tolerant to toxic, highly reactive, oxygen species, particularly superoxide anions, produced in the plant's cells as a result of a stress condition, particularly a naturally occurring stress condition, characterized by the step of: transforming the genome, preferably the nuclear genome, of said plant with the recombinant SOD gene of anyone of claims 1-3, whereby an SOD is produced, preferably overproduced, in cells of said plant.